

TFAP2C Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21677c

Product Information

Application	WB, E
Primary Accession	Q92754
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB48469
Calculated MW	49177

Additional Information

Gene ID	7022
Other Names	Transcription factor AP-2 gamma, AP2-gamma, Activating enhancer-binding protein 2 gamma, Transcription factor ERF-1, TFAP2C
Target/Specificity	This TFAP2C antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 114-147 amino acids from the central region of human TFAP2C.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TFAP2C Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TFAP2C
Function	Sequence-specific DNA-binding transcription factor that interacts with cellular enhancer elements to regulate transcription of selected genes, and which plays a key role in early embryonic development (PubMed: 11694877 , PubMed: 24413532). AP-2 factors bind to the consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of

important biological functions (PubMed:[11694877](#), PubMed:[24413532](#)). TFAP2C plays a key role in early embryonic development by regulating both inner cell mass (ICM) and trophoblast differentiation (By similarity). At the 8-cell stage, during morula development, controls expression of cell-polarity genes (By similarity). Upon trophoblast commitment, binds to late trophoblast genes in blastocysts together with CDX2, and later to extra-embryonic ectoderm genes together with SOX2 (By similarity). Binds to both closed and open chromatin with other transcription factors (By similarity). Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer (PubMed:[24413532](#)).

Cellular Location

Nucleus.


Background

Sequence-specific DNA-binding protein that interacts with inducible viral and cellular enhancer elements to regulate transcription of selected genes. AP-2 factors bind to the consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of important biological functions including proper eye, face, body wall, limb and neural tube development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha and MYC. Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer.

References

- Williamson J.A.,et al.Genomics 35:262-264(1996).
McPherson L.A.,et al.Proc. Natl. Acad. Sci. U.S.A. 94:4342-4347(1997).
Haselton M.D.,et al.Submitted (AUG-2001) to the EMBL/GenBank/DBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Deloukas P.,et al.Nature 414:865-871(2001).

Images



Anti-TFAP2C Antibody (Center) at 1:2000 dilution + SK-BR-3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 49 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.