

NDUFA7 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21706b

Product Information

Application	WB, E
Primary Accession	O95182
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53480
Calculated MW	12551

Additional Information

Gene ID	4701
Other Names	NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 7, Complex I-B145a, CI-B145a, NADH-ubiquinone oxidoreductase subunit B145a, NDUFA7
Target/Specificity	This NDUFA7 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 80-110 amino acids from human NDUFA7.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NDUFA7 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NDUFA7
Function	Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.

Cellular Location

Mitochondrion inner membrane; Peripheral membrane protein; Matrix side

Background

Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.

References


Loeffen J.L.C.M.,et al.Biochem. Biophys. Res. Commun. 253:415-422(1998).

Zhang Q.-H.,et al.Genome Res. 10:1546-1560(2000).

Murray J.,et al.J. Biol. Chem. 278:13619-13622(2003).

Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).

Images



All lanes : Anti-NDUFA7 Antibody (C-Term) at 1:2000 dilution Lane 1: mouse heart lysate Lane 2: MCF-7 whole cell lysate Lane 3: SW480 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 13 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.