

RAP2A Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21722b

Product Information

Application WB, E **Primary Accession** P10114 Reactivity Human Host Rabbit Clonality polyclonal Isotype Rabbit IgG **Clone Names** RB53398 **Calculated MW** 20615

Additional Information

Gene ID 5911

Other Names Ras-related protein Rap-2a, RbBP-30, RAP2A

Target/Specificity This RAP2A antibody is generated from a rabbit immunized with a KLH

conjugated synthetic peptide between 124-159 amino acids from human

RAP2A.

Dilution WB~~1:2000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions RAP2A Antibody (C-Term) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name RAP2A (HGNC:9861)

Function Small GTP-binding protein which cycles between a GDP-bound inactive and

a GTP-bound active form (PubMed: 14966141, PubMed: 15342639,

PubMed:16246175, PubMed:16540189, PubMed:18930710,

PubMed: 20159449, PubMed: 35293963). In its active form interacts with and

regulates several effectors including MAP4K4, MINK1 and TNIK (PubMed: 14966141, PubMed: 15342639, PubMed: 18930710,

PubMed:20159449). Part of a signaling complex composed of NEDD4, RAP2A and TNIK which regulates neuronal dendrite extension and arborization during development (PubMed:20159449). More generally, it is part of several signaling cascades and regulates cytoskeletal rearrangements, cell migration, cell adhesion and cell spreading (PubMed:14966141, PubMed:15342639, PubMed:16246175, PubMed:16540189, PubMed:18930710, PubMed:20159449, PubMed:35293963).

Cellular Location

Midbody. Cell projection, lamellipodium membrane. Golgi apparatus. Recycling endosome membrane; Lipid-anchor; Cytoplasmic side. Lysosome. Note=Localized to the Golgi (PubMed:35293963, PubMed:7962206). May also localize to the gelatinase- containing granules of neutrophils (PubMed:8391995). Colocalized with RASGEF1B to midbody at telophase (PubMed:23894443). Localized predominantly to the plasma membrane, where it is enriched at lamellipodia ruffles (PubMed:35293963). Cycles between the lamellipodia plasma membrane and endosomes when ubiquitinated by the ECS(RAB40B) complex (PubMed:35293963). Without the ubiquitin signal, sorted to lysosomes for degradation (PubMed:35293963)


Background

Small GTP-binding protein which cycles between a GDP- bound inactive and a GTP-bound active form. In its active form interacts with and regulates several effectors including MAP4K4, MINK1 and TNIK. Part of a signaling complex composed of NEDD4, RAP2A and TNIK which regulates neuronal dendrite extension and arborization during development. More generally, it is part of several signaling cascades and may regulate cytoskeletal rearrangements, cell migration, cell adhesion and cell spreading.

References

Pizon V., et al. Oncogene 3:201-204(1988). Fan Z.S., et al. Submitted (NOV-1999) to the EMBL/GenBank/DDBJ databases. Puhl H.L. III, et al. Submitted (MAR-2002) to the EMBL/GenBank/DDBJ databases. Ota T., et al. Nat. Genet. 36:40-45(2004). Dunham A., et al. Nature 428:522-528(2004).

Images



Anti-RAP2A Antibody (C-Term) at 1:2000 dilution + A431 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 21 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.