10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SAG Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21724b

Product Information

Application	WB, E
Primary Accession	P10523
Reactivity	Human, Rat, Mouse
Predicted	Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53546
Calculated MW	45120

Additional Information

Gene ID	6295
Other Names	S-arrestin, 48 kDa protein, Retinal S-antigen, S-AG, Rod photoreceptor arrestin, SAG
Target/Specificity	This SAG antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 273-307 amino acids from the human SAG.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SAG Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SAG
Function	Involved in the regulation of phototransduction, playing a key role in signal termination after rhodopsin (RHO) has been activated by light in rod photoreceptors. Binds to photoactivated and phosphorylated RHO, preventing its interaction with transducin (GNAT1) and terminating transducin-mediated

	opsin signaling (By similarity). May play a role in preventing light-dependent degeneration of retinal photoreceptor cells (PubMed:9565049).
Cellular Location	Photoreceptor inner segment {ECO:0000250|UniProtKB:P20443}. Cell projection, cilium, photoreceptor outer segment. Membrane {ECO:0000250|UniProtKB:P20443}; Peripheral membrane protein {ECO:0000250|UniProtKB:P20443}. Note=In dark-adapted retina, it is predominatly localized to the outer nuclear layer and photoreceptor inner segments (By similarity). Localized to photoreceptor outer segments in light-exposed retina. During light exposure it first accumulates in the proximal region of photoreceptor outer segments, and moves toward the distal tip of the outer segment as light exposure is prolonged (By similarity). {ECO:0000250|UniProtKB:P20443}
Tissue Location	Detected in retina, in the proximal portion of the outer segment of rod photoreceptor cells (at protein level)


Background

Arrestin is one of the major proteins of the ros (retinal rod outer segments); it binds to photoactivated-phosphorylated rhodopsin, thereby apparently preventing the transducin-mediated activation of phosphodiesterase.

References

Yamaki K.,et al.FEBS Lett. 234:39-43(1988).
Yamaki K.,et al.FEBS Lett. 236:507-507(1988).
Yamamoto S.,et al.Nat. Genet. 15:175-178(1997).
Hillier L.W.,et al.Nature 434:724-731(2005).
Roni V.,et al.BMC Genomics 8:42-42(2007).

Images



Anti-SAG Antibody (C-Term) at 1:2000 dilution + mouse eyeball lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 45 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.