

ADRB3 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21726c

Product Information

Application	WB, E
Primary Accession	P13945
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53477
Calculated MW	43519

Additional Information

Gene ID	155
Other Names	Beta-3 adrenergic receptor, Beta-3 adrenoreceptor, Beta-3 adrenoceptor, ADRB3, ADRB3R, B3AR
Target/Specificity	This ADRB3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 216-249 amino acids from the Central region of human ADRB3.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ADRB3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ADRB3
Synonyms	ADRB3R, B3AR
Function	Beta-adrenergic receptors mediate the catecholamine-induced activation of adenylate cyclase through the action of G proteins. Beta- 3 is involved in the regulation of lipolysis and thermogenesis.

Cellular Location Cell membrane; Multi-pass membrane protein.

Tissue Location Expressed mainly in adipose tissues.


Background

Beta-adrenergic receptors mediate the catecholamine- induced activation of adenylate cyclase through the action of G proteins. Beta-3 is involved in the regulation of lipolysis and thermogenesis.

References

Emorine L.J.,et al.Science 245:1118-1121(1989).
van Spronsen A.,et al.Eur. J. Biochem. 213:1117-1124(1993).
Lelias J.M.,et al.FEBS Lett. 324:127-130(1993).
Kopatz S.A.,et al.Submitted (NOV-2003) to the EMBL/GenBank/DDBJ databases.
Granneman J.G.,et al.Mol. Pharmacol. 42:964-970(1992).

Images



Anti-ADRB3 Antibody (Center) at 1:1000 dilution + HL-60 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 44 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.