

# RLBP1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP21727b

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P12271</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB49734
<b>Calculated MW</b>	36474

## Additional Information

---

<b>Gene ID</b>	6017
<b>Other Names</b>	Retinaldehyde-binding protein 1, Cellular retinaldehyde-binding protein, RLBP1, CRALBP
<b>Target/Specificity</b>	This RLBP1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 255-288 amino acids from the C-terminal region of human RLBP1.
<b>Dilution</b>	WB~~1:2000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	RLBP1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	RLBP1
<b>Synonyms</b>	CRALBP
<b>Function</b>	Soluble retinoid carrier essential the proper function of both rod and cone photoreceptors. Participates in the regeneration of active 11-cis-retinol and 11-cis-retinaldehyde, from the inactive 11- trans products of the rhodopsin

photocycle and in the de novo synthesis of these retinoids from 11-trans metabolic precursors. The cycling of retinoids between photoreceptor and adjacent pigment epithelium cells is known as the 'visual cycle'.

#### Cellular Location

Cytoplasm.

#### Tissue Location

Retina and pineal gland. Not present in photoreceptor cells but is expressed abundantly in the adjacent retinal pigment epithelium (RPE) and in the Mueller glial cells of the retina


## Background

Soluble retinoid carrier essential the proper function of both rod and cone photoreceptors. Participates in the regeneration of active 11-cis-retinol and 11-cis-retinaldehyde, from the inactive 11-trans products of the rhodopsin photocycle and in the de novo synthesis of these retinoids from 11-trans metabolic precursors. The cycling of retinoids between photoreceptor and adjacent pigment epithelium cells is known as the 'visual cycle'.

## References

Crabb J.W.,et al.J. Biol. Chem. 263:18688-18692(1988).  
Intres R.,et al.J. Biol. Chem. 269:25411-25418(1994).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.  
Crabb J.W.,et al.Protein Sci. 7:746-757(1998).

## Images



Anti-RLBP1 Antibody (C-term) at 1:2000 dilution + mouse eyeball lysate Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 36 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.