

CHRM4 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21733b

Product Information

Application	WB, E
Primary Accession	P08173
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53327
Calculated MW	53049

Additional Information

Gene ID	1132
Other Names	Muscarinic acetylcholine receptor M4, CHRM4
Target/Specificity	This CHRM4 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 201-236 amino acids from human CHRM4.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CHRM4 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CHRM4
Function	The muscarinic acetylcholine receptor mediates various cellular responses, including inhibition of adenylate cyclase, breakdown of phosphoinositides and modulation of potassium channels through the action of G proteins. Primary transducing effect is inhibition of adenylate cyclase.
Cellular Location	Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane;


Background

The muscarinic acetylcholine receptor mediates various cellular responses, including inhibition of adenylate cyclase, breakdown of phosphoinositides and modulation of potassium channels through the action of G proteins. Primary transducing effect is inhibition of adenylate cyclase.

References

Bonner T.I.,et al.Science 237:527-532(1987).
Bonner T.I.,et al.Neuron 1:403-410(1988).
Peralta E.G.,et al.EMBO J. 6:3923-3929(1987).
Puhl H.L. III,et al.Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Images



Anti-CHRM4 Antibody (C-Term) at 1:2000 dilution + human skeletal muscle lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 53 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.