10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

NDUFV1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21746c

Product Information

Application	WB, E
Primary Accession	P49821
Reactivity	Human, Mouse
Predicted	Rat, Bovine, Canine, Chicken
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53485
Calculated MW	50817

Additional Information

Gene ID	4723
Other Names	NADH dehydrogenase [ubiquinone] flavoprotein 1, mitochondrial, Complex I-51kD, CI-51kD, NADH dehydrogenase flavoprotein 1, NADH-ubiquinone oxidoreductase 51 kDa subunit, NDUFV1, UQOR1
Target/Specificity	This NDUFV1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 194-226 amino acids from the Central region of human NDUFV1.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NDUFV1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NDUFV1 (HGNC:7716)
Synonyms	UQOR1
Function	Core subunit of the mitochondrial membrane respiratory chain NADH

dehydrogenase (Complex I) which catalyzes electron transfer from NADH through the respiratory chain, using ubiquinone as an electron acceptor (PubMed:28844695). Part of the peripheral arm of the enzyme, where the electrons from NADH are accepted by flavin mononucleotide (FMN) and then passed along a chain of iron-sulfur clusters by electron tunnelling to the final acceptor ubiquinone (PubMed:28844695). Contains FMN, which is the initial electron acceptor as well as one iron-sulfur cluster (PubMed:28844695).

Cellular Location

Mitochondrion inner membrane {ECO:0000250|UniProtKB:P25708}; Peripheral membrane protein {ECO:0000250|UniProtKB:P25708}; Matrix side {ECO:0000250|UniProtKB:P25708}


Background

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone (By similarity).

References

de Coo R.F.M.,et al.Mamm. Genome 10:49-53(1999).
Schuelke M.,et al.Biochem. Biophys. Res. Commun. 245:599-606(1998).
Hu R.-M.,et al.Proc. Natl. Acad. Sci. U.S.A. 97:9543-9548(2000).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Images



All lanes : Anti-NDUFV1 Antibody (Center) at 1:2000 dilution Lane 1: mouse kidney lysate Lane 2: HepG2 whole cell lysate Lane 3: A431 whole cell lysate Lane 4: HL-60 whole cell lysate Lane 5: HT-29 whole cell lysate Lane 6: mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 51 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.