

ATXN2 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21752a

Product Information

Application	WB, E
Primary Accession	Q99700
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53600
Calculated MW	140283

Additional Information

Gene ID	6311
Other Names	Ataxin-2, Spinocerebellar ataxia type 2 protein, Trinucleotide repeat-containing gene 13 protein, ATXN2, ATX2, SCA2, TNRC13
Target/Specificity	This ATXN2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 321-355 amino acids from human ATXN2.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ATXN2 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ATXN2
Synonyms	ATX2, SCA2, TNRC13
Function	Involved in EGFR trafficking, acting as negative regulator of endocytic EGFR internalization at the plasma membrane.

Cellular Location	Cytoplasm.
Tissue Location	Expressed in the brain, heart, liver, skeletal muscle, pancreas and placenta. Isoform 1 is predominant in the brain and spinal cord. Isoform 4 is more abundant in the cerebellum. In the brain, broadly expressed in the amygdala, caudate nucleus, corpus callosum, hippocampus, hypothalamus, substantia nigra, subthalamic nucleus and thalamus.


Background

Involved in EGFR trafficking, acting as negative regulator of endocytic EGFR internalization at the plasma membrane.

References

Pulst S.-M.,et al.Nat. Genet. 14:269-276(1996).
Sanpei K.,et al.Nat. Genet. 14:277-284(1996).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Scherer S.E.,et al.Nature 440:346-351(2006).
Imbert G.,et al.Nat. Genet. 14:285-291(1996).

Images



All lanes : Anti-ATXN2 Antibody (N-Term) at 1:2000 dilution Lane 1: human brain lysate Lane 2: mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 140 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.