

WDR51B Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21756b

Product Information

Application	WB, E
Primary Accession	Q8TC44
Reactivity	Human
Predicted	Canine
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB47250
Calculated MW	53668

Additional Information

Gene ID	282809
Other Names	POC1 centriolar protein homolog B, Pix1, Proteome of centriole protein 1B, WD repeat-containing protein 51B, POC1B, WDR51B
Target/Specificity	This WDR51B antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 386-417 amino acids from the C-terminal region of human WDR51B.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	WDR51B Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	POC1B (HGNC:30836)
Synonyms	WDR51B
Function	Plays an important role in centriole assembly and/or stability and ciliogenesis (PubMed: 20008567 , PubMed: 32060285). Involved in early steps of

centriole duplication, as well as in the later steps of centriole length control (PubMed:[19109428](#)). Acts in concert with POC1A to ensure centriole integrity and proper mitotic spindle formation (PubMed:[32060285](#)). Required for primary cilia formation, ciliary length and also cell proliferation (PubMed:[23015594](#)). Required for retinal integrity (PubMed:[25044745](#)). Acts as a positive regulator of centriole elongation (PubMed:[37934472](#)).

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Cytoplasm, cytoskeleton, cilium basal body Cytoplasm, cytoskeleton, spindle pole. Note=Component of both mother and daughter centrioles (PubMed:32060285). Localizes to the basal body and centriole adjacent to the connecting cilium of photoreceptors and in synapses of the outer plexiform layer. Localizes to the inner scaffold in the central region of centrioles {ECO:0000250 | UniProtKB:Q8BHD1, ECO:0000269 | PubMed:32060285, ECO:0000269 | PubMed:37934472}

Tissue Location

Expressed in the retina.


Background

Plays an important role in centriole assembly and/or stability and ciliogenesis (PubMed: [20008567](#)). Involved in early steps of centriole duplication, as well as in the later steps of centriole length control (PubMed:[19109428](#)). Acts in concert with POC1A to ensure centriole integrity and proper mitotic spindle formation. Required for primary cilia formation, ciliary length and also cell proliferation (PubMed:[23015594](#)). Required for retinal integrity (PubMed:[25044745](#)).

References

Ota T.,et al.Nat. Genet. 36:40-45(2004).
Scherer S.E.,et al.Nature 440:346-351(2006).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Hames R.S.,et al.Exp. Cell Res. 314:574-589(2008).
Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).

Images



All lanes : Anti-WDR51B Antibody (C-term) at 1:2000 dilution
Lane 1: human testis lysate
Lane 2: MDA-MB-453 whole cell lysate
Lane 3: TT whole cell lysate
Lane 4: Y79 whole cell lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 84 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.