

KAT5 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21760b

Product Information

Application	WB, E
Primary Accession	Q92993
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53893
Calculated MW	58582

Additional Information

Gene ID	10524
Other Names	Histone acetyltransferase KAT5, 60 kDa Tat-interactive protein, Tip60, Histone acetyltransferase HTATIP, HIV-1 Tat interactive protein, Lysine acetyltransferase 5, cPLA(2)-interacting protein, KAT5, HTATIP, TIP60
Target/Specificity	This KAT5 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 367-398 amino acids from human KAT5.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	KAT5 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	KAT5 {ECO:0000303 PubMed:32817552, ECO:0000312 HGNC:HGNC:5275}
Function	Catalytic subunit of the NuA4 histone acetyltransferase complex, a multiprotein complex involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H2A and H4 (PubMed: 12776177 , PubMed: 14966270 , PubMed: 15042092 ,

PubMed:[15121871](#), PubMed:[15310756](#), PubMed:[16387653](#), PubMed:[19909775](#), PubMed:[25865756](#), PubMed:[27153538](#), PubMed:[29174981](#), PubMed:[29335245](#), PubMed:[32822602](#), PubMed:[33076429](#)). Histone acetylation alters nucleosome-DNA interactions and promotes interaction of the modified histones with other proteins which positively regulate transcription (PubMed:[12776177](#), PubMed:[14966270](#), PubMed:[15042092](#), PubMed:[15121871](#), PubMed:[15310756](#)). The NuA4 histone acetyltransferase complex is required for the activation of transcriptional programs associated with proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair (PubMed:[17709392](#), PubMed:[19783983](#), PubMed:[32832608](#)). The NuA4 complex plays a direct role in repair of DNA double-strand breaks (DSBs) by promoting homologous recombination (HR): the complex inhibits TP53BP1 binding to chromatin via MBTD1, which recognizes and binds histone H4 trimethylated at 'Lys-20' (H4K20me), and KAT5 that catalyzes acetylation of 'Lys-15' of histone H2A (H2AK15ac), thereby blocking the ubiquitination mark required for TP53BP1 localization at DNA breaks (PubMed:[27153538](#), PubMed:[32832608](#)). Also involved in DSB repair by mediating acetylation of 'Lys-5' of histone H2AX (H2AXK5ac), promoting NBN/NBS1 assembly at the sites of DNA damage (PubMed:[17709392](#), PubMed:[26438602](#)). The NuA4 complex plays a key role in hematopoietic stem cell maintenance and is required to maintain acetylated H2A.Z/H2AZ1 at MYC target genes (By similarity). The NuA4 complex is also required for spermatid development by promoting acetylation of histones: histone hyperacetylation is required for histone replacement during the transition from round to elongating spermatids (By similarity). Component of a SWR1-like complex that specifically mediates the removal of histone H2A.Z/H2AZ1 from the nucleosome (PubMed:[24463511](#)). Also acetylates non-histone proteins, such as BMAL1, ATM, AURKB, CHKA, CGAS, ERCC4/XPF, LPIN1, TP53/p53, NDC80/HEC1, NR1D2, RAN, SOX4, FOXP3, SQSTM1, ULK1 and RUBCNL/Pacer (PubMed:[16141325](#), PubMed:[17189187](#), PubMed:[17360565](#), PubMed:[17996965](#), PubMed:[24835996](#), PubMed:[26829474](#), PubMed:[29040603](#), PubMed:[30409912](#), PubMed:[30704899](#), PubMed:[31857589](#), PubMed:[32034146](#), PubMed:[32817552](#), PubMed:[34077757](#)). Directly acetylates and activates ATM (PubMed:[16141325](#)). Promotes nucleotide excision repair (NER) by mediating acetylation of ERCC4/XPF, thereby promoting formation of the ERCC4-ERCC1 complex (PubMed:[32034146](#)). Relieves NR1D2-mediated inhibition of APOC3 expression by acetylating NR1D2 (PubMed:[17996965](#)). Acts as a regulator of regulatory T-cells (Treg) by catalyzing FOXP3 acetylation, thereby promoting FOXP3 transcriptional repressor activity (PubMed:[17360565](#), PubMed:[24835996](#)). Involved in skeletal myoblast differentiation by mediating acetylation of SOX4 (PubMed:[26291311](#)). Catalyzes acetylation of APBB1/FE65, increasing its transcription activator activity (PubMed:[33938178](#)). Promotes transcription elongation during the activation phase of the circadian cycle by catalyzing acetylation of BMAL1, promoting elongation of circadian transcripts (By similarity). Together with GSK3 (GSK3A or GSK3B), acts as a regulator of autophagy: phosphorylated at Ser-86 by GSK3 under starvation conditions, leading to activate acetyltransferase activity and promote acetylation of key autophagy regulators, such as ULK1 and RUBCNL/Pacer (PubMed:[30704899](#)). Acts as a regulator of the cGAS-STING innate antiviral response by catalyzing acetylation the N-terminus of CGAS, thereby promoting CGAS DNA-binding and activation (PubMed:[32817552](#)). Also regulates lipid metabolism by mediating acetylation of CHKA or LPIN1 (PubMed:[34077757](#)). Promotes lipolysis of lipid droplets following glucose deprivation by mediating acetylation of isoform 1 of CHKA, thereby promoting monomerization of CHKA and its conversion into a tyrosine-protein kinase (PubMed:[34077757](#)). Acts as a regulator of fatty-acid-induced triacylglycerol synthesis by catalyzing acetylation of LPIN1, thereby promoting the synthesis of diacylglycerol (PubMed:[29765047](#)). In addition to protein acetyltransferase, can use

different acyl-CoA substrates, such as (2E)-butenoyl-CoA (crotonyl-CoA), S-lactoyl-CoA (lactyl-CoA) and 2-hydroxyisobutanoyl-CoA (2-hydroxyisobutyryl-CoA), and is able to mediate protein crotonylation, lactylation and 2-hydroxyisobutyrylation, respectively (PubMed:[29192674](#), PubMed:[34608293](#), PubMed:[38961290](#)). Acts as a key regulator of chromosome segregation and kinetochore-microtubule attachment during mitosis by mediating acetylation or crotonylation of target proteins (PubMed:[26829474](#), PubMed:[29040603](#), PubMed:[30409912](#), PubMed:[34608293](#)). Catalyzes acetylation of AURKB at kinetochores, increasing AURKB activity and promoting accurate chromosome segregation in mitosis (PubMed:[26829474](#)). Acetylates RAN during mitosis, promoting microtubule assembly at mitotic chromosomes (PubMed:[29040603](#)). Acetylates NDC80/HEC1 during mitosis, promoting robust kinetochore-microtubule attachment (PubMed:[30409912](#)). Catalyzes crotonylation of MAPRE1/EB1, thereby ensuring accurate spindle positioning in mitosis (PubMed:[34608293](#)). Catalyzes lactylation of NBN/NBS1 in response to DNA damage, thereby promoting DNA double-strand breaks (DSBs) via homologous recombination (HR) (PubMed:[38961290](#)).

Cellular Location

Nucleus. Chromosome. Cytoplasm Chromosome, centromere, kinetochore Cytoplasm, cytoskeleton, spindle pole Nucleus, nucleolus. Cytoplasm, perinuclear region. Note=Upon stimulation with EDN1, it is exported from the nucleus to the perinuclear region and UV irradiation induces translocation into punctuate subnuclear structures named nuclear bodies (PubMed:[11262386](#)). Transiently localizes to kinetochores in early mitosis (PubMed:[26829474](#)). Localizes to spindle poles when chromosomes align during metaphase (PubMed:[34608293](#)). Localizes in the cytoplasm and nucleus of round spermatids (By similarity). {ECO:0000250|UniProtKB:Q8CHK4, ECO:0000269|PubMed:[11262386](#), ECO:0000269|PubMed:[26829474](#), ECO:0000269|PubMed:[34608293](#)}


Background

Catalytic subunit of the NuA4 histone acetyltransferase complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome-DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. NuA4 may also play a direct role in DNA repair when recruited to sites of DNA damage. Directly acetylates and activates ATM. Component of a SWR1-like complex that specifically mediates the removal of histone H2A.Z/H2AFZ from the nucleosome. In case of HIV-1 infection, interaction with the viral Tat protein leads to KAT5 polyubiquitination and targets it to degradation. Relieves NR1D2-mediated inhibition of APOC3 expression by acetylating NR1D2.

References

Kamine J.,et al.Virology 216:357-366(1996).
 Sheridan A.M.,et al.Mol. Cell. Biol. 21:4470-4481(2001).
 Legube G.,et al.Gene 310:161-168(2003).
 Ota T.,et al.Nat. Genet. 36:40-45(2004).
 Taylor T.D.,et al.Nature 440:497-500(2006).

Images



All lanes : Anti-KAT5 Antibody (C-Term) at 1:2000 dilution
Lane 1: PC-3 whole cell lysate Lane 2: SK-BR-3 whole cell lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 59 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.