

BMP4 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21762a

Product Information

Application	WB, E
Primary Accession	<u>P12644</u>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53905
Calculated MW	46555

Additional Information

Gene ID	652
Other Names	Bone morphogenetic protein 4, BMP-4, Bone morphogenetic protein 2B, BMP-2B, BMP4, BMP2B, DVR4
Target/Specificity	This BMP4 antibody is generated from a rabbit immunized with a recombinant protein of human BMP4.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	BMP4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	BMP4 (<u>HGNC:1071</u>)
Function	Growth factor of the TGF-beta superfamily that plays essential roles in many developmental processes, including neurogenesis, vascular development, angiogenesis and osteogenesis (PubMed: <u>31363885</u>). Acts in concert with PTHLH/PTHRP to stimulate ductal outgrowth during embryonic mammary development and to inhibit hair follicle induction (By similarity). Initiates the canonical BMP signaling cascade by associating with type I receptor BMPR1A

	and type II receptor BMPR2 (PubMed: <u>25868050</u> , PubMed: <u>8006002</u>). Once all three components are bound together in a complex at the cell surface, BMPR2 phosphorylates and activates BMPR1A. In turn, BMPR1A propagates signal by phosphorylating SMAD1/5/8 that travel to the nucleus and act as activators and repressors of transcription of target genes (PubMed: <u>25868050</u> , PubMed: <u>29212066</u>). Positively regulates the expression of odontogenic development regulator MSX1 via inducing the IPO7- mediated import of SMAD1 to the nucleus (By similarity). Required for MSX1-mediated mesenchymal molar tooth bud development beyond the bud stage, via promoting Wnt signaling (By similarity). Acts as a positive regulator of odontoblast differentiation during mesenchymal tooth germ formation, expression is repressed during the bell stage by MSX1- mediated inhibition of CTNNB1 signaling (By similarity). Able to induce its own expression in dental mesenchymal cells and also in the neighboring dental epithelial cells via an MSX1-mediated pathway (By similarity). Can also signal through non-canonical BMP pathways such as ERK/MAP kinase, PI3K/Akt, or SRC cascades (PubMed: <u>31363885</u>). For example, induces SRC phosphorylation which, in turn, activates VEGFR2, leading to an angiogenic response (PubMed: <u>31363885</u>).
Cellular Location	Secreted, extracellular space, extracellular matrix
Tissue Location	Expressed in the lung and lower levels seen in the kidney. Present also in normal and neoplastic prostate tissues, and prostate cancer cell lines

Background

Induces cartilage and bone formation. Also act in mesoderm induction, tooth development, limb formation and fracture repair. Acts in concert with PTHLH/PTHRP to stimulate ductal outgrowth during embryonic mammary development and to inhibit hair follicle induction (By similarity).

References

Wozney J.M.,et al.Science 242:1528-1534(1988). Shore E.M.,et al.Calcif. Tissue Int. 63:221-229(1998). Oida S.,et al.DNA Seq. 5:273-275(1995). Yanagita M.,et al.Biochem. Biophys. Res. Commun. 316:490-500(2004). Felder B.,et al.Eur. J. Hum. Genet. 10:753-756(2002).

Images



All lanes : Anti-BMP4 Antibody at 1:1000-1:2000 dilution Lane 1: A549 whole cell lysate Lane 2: human brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 47 kDa Blocking/Dilution buffer: 5% NFDM/TBST. Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.