10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

FBXO33 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21765b

Product Information

Application	WB, E
Primary Accession	Q7Z6M2
Reactivity	Human, Mouse
Predicted	Rat, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53625
Calculated MW	62630

Additional Information

Gene ID	254170
Other Names	F-box only protein 33, FBXO33, FBX33
Target/Specificity	This FBXO33 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 416-449 amino acids from human FBXO33.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	FBXO33 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	FBXO33
Synonyms	FBX33
Function	Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins. Probably recognizes

and binds to phosphorylated target proteins. Recognizes YBX1 (By similarity).


Background

Substrate recognition component of a SCF (SKP1-CUL1-F- box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins. Probably recognizes and binds to phosphorylated target proteins. Recognizes YBX1 (By similarity).

References

Li W.B.,et al.Submitted (FEB-2003) to the EMBL/GenBank/DDBJ databases.

Images



Anti-FBXO33 Antibody (C-Term) at 1:2000 dilution + mouse testis lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 63 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.