

BAG3 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21826a

Product Information

Application	WB, E
Primary Accession	O95817
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB54138
Calculated MW	61595

Additional Information

Gene ID	9531
Other Names	BAG family molecular chaperone regulator 3, BAG-3, Bcl-2-associated athanogene 3, Bcl-2-binding protein Bis, Docking protein CAIR-1, BAG3, BIS
Target/Specificity	This BAG3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 31-62 amino acids from human BAG3.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	BAG3 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	BAG3
Synonyms	BIS
Function	Co-chaperone and adapter protein that connects different classes of molecular chaperones including heat shock proteins 70 (HSP70s), e.g. HSPA1A/HSP70 or HSPA8/HSC70, and small heat shock proteins (sHSPs), e.g. HSPB8 (PubMed: 27884606 , PubMed: 30559338). Acts as a

nucleotide-exchange factor (NEF) promoting the release of ADP from HSP70s, thereby triggering client protein release (PubMed:[27884606](#), PubMed:[30559338](#)). Nucleotide release is mediated via BAG3 binding to the nucleotide-binding domain (NBD) of HSP70s, whereas client release is mediated via binding to the substrate-binding domain (SBD) (PubMed:[27474739](#), PubMed:[9873016](#)). Has anti-apoptotic activity (PubMed:[10597216](#)). Plays a role in the HSF1 nucleocytoplasmic transport (PubMed:[26159920](#)).

Cellular Location

Nucleus. Cytoplasm. Note=Colocalizes with HSF1 to the nucleus upon heat stress (PubMed:[26159920](#))


Background

Inhibits the chaperone activity of HSP70/HSC70 by promoting substrate release. Has anti-apoptotic activity.

References

Takayama S.,et al.J. Biol. Chem. 274:781-786(1999).
Lee J.H.,et al.Oncogene 18:6183-6190(1999).
Doong H.,et al.Oncogene 19:4385-4395(2000).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).

Images



All lanes : Anti-BAG3 Antibody (N-Term) at 1:2000 dilution
Lane 1: A431 whole cell lysate Lane 2: MCF-7 whole cell lysate Lane 3: U-2OS whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 62 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.