

# PYGL Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21847b

## Product Information

---

<b>Application</b>	WB, IHC-P, E
<b>Primary Accession</b>	<a href="#">P06737</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB53470
<b>Calculated MW</b>	97149

## Additional Information

---

<b>Gene ID</b>	5836
<b>Other Names</b>	Glycogen phosphorylase, liver form, PYGL
<b>Target/Specificity</b>	This PYGL antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 809-843 amino acids from human PYGL.
<b>Dilution</b>	WB~~1:2000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	PYGL Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	PYGL ( <a href="#">HGNC:9725</a> )
<b>Function</b>	Allosteric enzyme that catalyzes the rate-limiting step in glycogen catabolism, the phosphorolytic cleavage of glycogen to produce glucose-1-phosphate, and plays a central role in maintaining cellular and organismal glucose homeostasis.
<b>Cellular Location</b>	Cytoplasm, cytosol.


## Background

Phosphorylase is an important allosteric enzyme in carbohydrate metabolism. Enzymes from different sources differ in their regulatory mechanisms and in their natural substrates. However, all known phosphorylases share catalytic and structural properties.


## References

Newgard C.B.,et al.Proc. Natl. Acad. Sci. U.S.A. 83:8132-8136(1986).  
Carty M.D.,et al.Submitted (MAY-1998) to the EMBL/GenBank/DDBJ databases.  
Chang S.,et al.Hum. Mol. Genet. 7:865-870(1998).  
Burwinkel B.,et al.Am. J. Hum. Genet. 62:785-791(1998).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).

## Images



All lanes : Anti-PYGL Antibody (C-Term) at 1:2000 dilution  
Lane 1: human liver lysate Lane 2: HeLa whole cell lysate  
Lane 3: HepG2 whole cell lysate Lane 4: NIH/3T3 whole cell lysate Lane 5: PC-3 whole cell lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 97 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.



AP21847b staining PYGL in human liver tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.