10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MYH4 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21870a

Product Information

Application	WB, E
Primary Accession	Q9Y623
Other Accession	Q076A3, Q9UKX3, Q9BE40, Q076A6, Q8MJV0, Q5SX40, Q9TV61, P02565, Q076A5, Q9TV62, Q28641, Q29RW1, P13539, Q02566, P02563, Q9BE39, P49824, Q8MJU9, P79293, P13538, Q90339
Reactivity	Human
Predicted	Bovine, Chicken, Human, Mouse, Pig, Rat, Horse, Canine, Rabbit
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53412
Calculated MW	223071

Additional Information

Gene ID	4622
Other Names	Myosin-4, Myosin heavy chain 2b, MyHC-2b, Myosin heavy chain 4, Myosin heavy chain IIb, MyHC-IIb, Myosin heavy chain, skeletal muscle, fetal, MYH4
Target/Specificity	This MYH4 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 482-515 amino acids from human MYH4.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MYH4 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MYH4
Function	Muscle contraction.

Cellular Location Cytoplasm, myofibril. Note=Thick filaments of the myofibrils


Background

Muscle contraction.

References

Weiss A.,et al.J. Mol. Biol. 290:61-75(1999).
Zody M.C.,et al.Nature 440:1045-1049(2006).
Denis N.J.,et al.Proteomics 7:868-874(2007).

Images



Anti-MYH4 Antibody (N-Term) at 1:2000 dilution + human skeletal muscle lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 223 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.