

# FANCM Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP21872b

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">Q8IYD8</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB54125
<b>Calculated MW</b>	232191

## Additional Information

---

<b>Gene ID</b>	57697
<b>Other Names</b>	Fanconi anemia group M protein, Protein FACM, ATP-dependent RNA helicase FANCM, Fanconi anemia-associated polypeptide of 250 kDa, FAAP250, Protein Hef ortholog, FANCM, KIAA1596
<b>Target/Specificity</b>	This FANCM antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1752-1783 amino acids from human FANCM.
<b>Dilution</b>	WB~~1:2000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	FANCM Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	FANCM
<b>Synonyms</b>	KIAA1596
<b>Function</b>	DNA-dependent ATPase component of the Fanconi anemia (FA) core complex (PubMed: <a href="#">16116422</a> ). Required for the normal activation of the FA

pathway, leading to monoubiquitination of the FANCI-FANCD2 complex in response to DNA damage, cellular resistance to DNA cross-linking drugs, and prevention of chromosomal breakage (PubMed:16116422, PubMed:19423727, PubMed:20347428, PubMed:20347429, PubMed:29231814). In complex with CENPS and CENPX, binds double-stranded DNA (dsDNA), fork-structured DNA (fsDNA) and Holliday junction substrates (PubMed:20347428, PubMed:20347429). Its ATP-dependent DNA branch migration activity can process branched DNA structures such as a movable replication fork. This activity is strongly stimulated in the presence of CENPS and CENPX (PubMed:20347429). In complex with FAAP24, efficiently binds to single-strand DNA (ssDNA), splayed-arm DNA, and 3'-flap substrates (PubMed:17289582). In vitro, on its own, strongly binds ssDNA oligomers and weakly fsDNA, but does not bind to dsDNA (PubMed:16116434).

**Cellular Location**

Nucleus

**Tissue Location**

Expressed in germ cells of fetal and adult ovaries. In fetal ovaries, it is present in oogonia but expression is stronger in pachytene stage oocytes. Expressed in oocytes arrested at the diplotene stage of prophase I during the last trimester of pregnancy and in adults (PubMed:29231814). Expressed in the testis (PubMed:30075111).


## Background

ATPase required for FANCD2 ubiquitination, a key reaction in DNA repair. Binds to ssDNA but not to dsDNA. Recruited to forks stalled by DNA interstrand cross-links, and required for cellular resistance to such lesions.

## References

Meetei A.R., et al. Nat. Genet. 37:958-963(2005).  
Ota T., et al. Nat. Genet. 36:40-45(2004).  
Heilig R., et al. Nature 421:601-607(2003).  
Nagase T., et al. DNA Res. 7:273-281(2000).  
Mosedale G., et al. Nat. Struct. Mol. Biol. 12:763-771(2005).

## Images



All lanes : Anti-FANCM Antibody (C-Term) at 1:2000 dilution  
Lane 1: HeLa whole cell lysate  
Lane 2: U-2OS whole cell lysate  
Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 232 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.