

TMEM184C Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21875b

Product Information

Application	WB, E
Primary Accession	Q9NVA4
Other Accession	Q5RET6
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB54282
Calculated MW	50142

Additional Information

Gene ID	55751
Other Names	Transmembrane protein 184C, Transmembrane protein 34, TMEM184C, TMEM34
Target/Specificity	This TMEM184C antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 407-438 amino acids from human TMEM184C.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TMEM184C Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TMEM184C
Synonyms	TMEM34
Function	Possible tumor suppressor which may play a role in cell growth.

Cellular Location	Membrane; Multi-pass membrane protein
Tissue Location	Widely expressed with higher expression in lung, kidney, spleen, pancreas, thymus, prostate, testis, ovary, small intestine and thyroid.


Background

Possible tumor suppressor which may play a role in cell growth.

References

Ota T.,et al.Nat. Genet. 36:40-45(2004).
 Hillier L.W.,et al.Nature 434:724-731(2005).
 Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
 Yu Y.,et al.Genome Res. 11:1392-1403(2001).
 Akaishi J.,et al.J. Cancer Res. Clin. Oncol. 133:213-218(2007).

Images



All lanes : Anti-TMEM184C Antibody (C-Term) at 1:2000 dilution
 Lane 1: HT-1080 whole cell lysate
 Lane 2: MCF-7 whole cell lysate
 Lysates/proteins at 20 µg per lane.
 Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 50 kDa
 Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.