

NPRL3 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21877a

Product Information

Application	WB, E
Primary Accession	Q12980
Reactivity	Human, Mouse
Predicted	Rat, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB54296
Calculated MW	63605

Additional Information

Gene ID	8131
Other Names	Nitrogen permease regulator 3-like protein, -14 gene protein, Alpha-globin regulatory element-containing gene protein, Protein CGTHBA, NPRL3, C16orf35, CGTHBA, MARE
Target/Specificity	This NPRL3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 86-117 amino acids from human NPRL3.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NPRL3 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NPRL3 {ECO:0000303 PubMed:26505888, ECO:0000312 HGNC:HGNC:14124}
Function	As a component of the GATOR1 complex functions as an inhibitor of the amino acid-sensing branch of the mTORC1 pathway (PubMed: 23723238 ,

PubMed:[29590090](#), PubMed:[35338845](#)). In response to amino acid depletion, the GATOR1 complex has GTPase activating protein (GAP) activity and strongly increases GTP hydrolysis by RagA/RRAGA (or RagB/RRAGB) within heterodimeric Rag complexes, thereby turning them into their inactive GDP-bound form, releasing mTORC1 from lysosomal surface and inhibiting mTORC1 signaling (PubMed:[23723238](#), PubMed:[29590090](#), PubMed:[35338845](#)). In the presence of abundant amino acids, the GATOR1 complex is negatively regulated by GATOR2, the other GATOR subcomplex, in this amino acid-sensing branch of the TORC1 pathway (PubMed:[23723238](#)).

Cellular Location

Lysosome membrane. Note=Localization to lysosomes is mediated by the KICSTOR complex and is amino acid-independent.

Tissue Location

Widely expressed. Expressed in the frontal lobe cortex as well as in the temporal, parietal, and occipital lobes (PubMed:26505888, PubMed:27173016).


Background

As a component of the GATOR1 complex, inhibitor of the amino acid-sensing branch of the TORC1 pathway. The GATOR1 complex strongly increases GTP hydrolysis by RRAGA and RRAGB within RRAGC-containing heterodimers, thereby deactivating RRGAs, releasing mTORC1 from lysosomal surface and inhibiting mTORC1 signaling.

References

Vyas P.,et al.Genomics 29:679-689(1995).
De Gobbi M.,et al.Science 312:1215-1217(2006).
Martin J.,et al.Nature 432:988-994(2004).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Neklesa T.K.,et al.PLoS Genet. 5:E1000515-E1000515(2009).

Images



Anti-NPRL3 Antibody (N-Term) at 1:2000 dilution + HT-1080 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 64 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.