

PICALM Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21881b

Product Information

Application	WB, E
Primary Accession	Q13492
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB54131
Calculated MW	70755

Additional Information

Gene ID	8301
Other Names	Phosphatidylinositol-binding clathrin assembly protein, Clathrin assembly lymphoid myeloid leukemia protein, PICALM, CALM
Target/Specificity	This PICALM antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 512-543 amino acids from human PICALM.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PICALM Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PICALM
Synonyms	CALM
Function	Cytoplasmic adapter protein that plays a critical role in clathrin-mediated endocytosis which is important in processes such as internalization of cell receptors, synaptic transmission or removal of apoptotic cells. Recruits AP-2

and attaches clathrin triskelions to the cytoplasmic side of plasma membrane leading to clathrin-coated vesicles (CCVs) assembly (PubMed:[10436022](#), PubMed:[16262731](#), PubMed:[27574975](#)). Furthermore, regulates clathrin-coated vesicle size and maturation by directly sensing and driving membrane curvature (PubMed:[25898166](#)). In addition to binding to clathrin, mediates the endocytosis of small R- SNAREs (Soluble NSF Attachment Protein REceptors) between plasma membranes and endosomes including VAMP2, VAMP3, VAMP4, VAMP7 or VAMP8 (PubMed:[21808019](#), PubMed:[22118466](#), PubMed:[23741335](#)). In turn, PICALM- dependent SNARE endocytosis is required for the formation and maturation of autophagic precursors (PubMed:[25241929](#)). Modulates thereby autophagy and the turnover of autophagy substrates such as MAPT/TAU or amyloid precursor protein cleaved C-terminal fragment (APP- CTF) (PubMed:[24067654](#), PubMed:[25241929](#)).

Cellular Location

Cell membrane. Membrane, clathrin-coated pit. Golgi apparatus. Cytoplasmic vesicle, clathrin- coated vesicle. Nucleus. Note=Colocalized with clathrin in the Golgi area (PubMed:10436022). Interaction with PIMREG may target PICALM to the nucleus in some cells (PubMed:16491119)

Tissue Location

Expressed in all tissues examined.


Background

Assembly protein recruiting clathrin and adapter protein complex 2 (AP2) to cell membranes at sites of coated-pit formation and clathrin-vesicle assembly. May be required to determine the amount of membrane to be recycled, possibly by regulating the size of the clathrin cage. Involved in AP2-dependent clathrin-mediated endocytosis at the neuromuscular junction.

References

Dreyling M.H.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:4804-4809(1996).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Nakajima D.,et al.Submitted (MAR-2005) to the EMBL/GenBank/DDBJ databases.
Taylor T.D.,et al.Nature 440:497-500(2006).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Images



Anti-PICALM Antibody (C-Term) at 1:2000 dilution + HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 71 kDa Blocking/Dilution buffer: 5% NFDM/TBST.