

# RNF126 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP21884c

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">Q9BV68</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB54136
<b>Calculated MW</b>	33861

## Additional Information

---

<b>Gene ID</b>	55658
<b>Other Names</b>	E3 ubiquitin-protein ligase RNF126, 632-, RING finger protein 126 {ECO:0000312 HGNC:HGNC:21151}, RNF126 ( <a href="http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=21151" target="_blank">HGNC:21151</a> )
<b>Target/Specificity</b>	This RNF126 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 84-115 amino acids from the Central region of human RNF126.
<b>Dilution</b>	WB~1:2000
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	RNF126 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	RNF126 ( <a href="#">HGNC:21151</a> )
<b>Function</b>	E3 ubiquitin-protein ligase that mediates ubiquitination of target proteins (PubMed: <a href="#">23277564</a> , PubMed: <a href="#">24275455</a> , PubMed: <a href="#">24981174</a> ). Depending on the associated E2 ligase, mediates 'Lys-48'- and 'Lys-63'- linked

polyubiquitination of substrates (By similarity). Part of a BAG6-dependent quality control process ensuring that proteins of the secretory pathway that are mislocalized to the cytosol are degraded by the proteasome. Probably acts by providing the ubiquitin ligase activity associated with the BAG6 complex and be responsible for ubiquitination of the hydrophobic mislocalized proteins and their targeting to the proteasome (PubMed:[24981174](#), PubMed:[29042515](#)). May also play a role in the endosomal recycling of IGF2R, the cation- independent mannose-6-phosphate receptor (PubMed:[24275455](#)). May play a role in the endosomal sorting and degradation of several membrane receptors including EGFR, FLT3, MET and CXCR4, by mediating their ubiquitination (PubMed:[23418353](#)). By ubiquitinating CDKN1A/p21 and targeting it for degradation, may also promote cell proliferation (PubMed:[23026136](#)). May monoubiquitinate AICDA (PubMed:[23277564](#)).

**Cellular Location** Cytoplasm. Nucleus

**Tissue Location** Highly expressed in liver and testis.


## Background

E3 ubiquitin-protein ligase that regulates several biological processes through ubiquitination of various target proteins. Depending on the associated E2 ligase, mediates 'Lys- 48'- and 'Lys-63'-linked polyubiquitination of substrates. Through their polyubiquitination, may play a role in the endosomal sorting and degradation of several membrane receptors including EGFR, FLT3, MET and CXCR4. May also be part of a BAG6-dependent quality control process ensuring that proteins of the secretory pathway that are mislocalized to the cytosol are degraded by the proteasome. May provide the ubiquitin ligase activity associated with the BAG6 complex and be responsible for ubiquitination of the mislocalized proteins and their targeting to the proteasome (PubMed:[24981174](#)). May also play a role in the endosomal recycling of IGF2R, the cation-independent mannose-6-phosphate receptor (PubMed:[24275455](#)). By ubiquitinating CDKN1A/p21 and targeting it for degradation, may also promote cell proliferation (PubMed:[23026136](#)). May monoubiquitinate AICDA (PubMed:[23277564](#)).

## References

- Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Daub H.,et al.Mol. Cell 31:438-448(2008).  
Olsen J.V.,et al.Sci. Signal. 3:RA3-RA3(2010).  
Zhi X.,et al.Cancer Res. 73:385-394(2013).  
Smith C.J.,et al.J. Cell Sci. 126:1366-1380(2013).

## Images



Anti-RNF126 Antibody (Center) at 1:2000 dilution + CEM whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 36 kDa Blocking/Dilution buffer: 5% NFDM/TBST.