10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

GNG2 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21903a

Product Information

Application	WB, IHC-P, E
Primary Accession	P59768
Other Accession	P63212, P63213, Q5R7U4
Reactivity	Human, Mouse
Predicted	Mouse, Bovine, Rat, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB54059
Calculated MW	7850

Additional Information

Gene ID	54331
Other Names	Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2, G gamma-I, GNG2
Target/Specificity	This GNG2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 19-52 amino acids from of human GNG2.
Dilution	WB~~1:8000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GNG2 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GNG2
Function	Guanine nucleotide-binding proteins (G proteins) are involved as a modulator or transducer in various transmembrane signaling systems (PubMed:29925951, PubMed:33762731, PubMed:34239069,

PubMed:35610220, PubMed:35714614, PubMed:35835867, PubMed:36087581, PubMed:36989299, PubMed:37327704, PubMed:37935376, PubMed:37935377, PubMed:37963465, PubMed:38168118, PubMed:38552625, PubMed:39103320). The beta and gamma chains are required for the GTPase activity, for replacement of GDP by GTP, and for G protein-effector interaction (PubMed:29925951, PubMed:33762731, PubMed:34239069, PubMed:35610220, PubMed:35714614, PubMed:35835867, PubMed:36087581, PubMed:36989299, PubMed:37327704, PubMed:37935376, PubMed:37935377, PubMed:37963465, PubMed:38168118, PubMed:38552625, PubMed:39103320). Involved in bitter and sweet taste transduction as well as in umami (monosodium glutamate, monopotassium glutamate, and inosine monophosphate) taste transduction (PubMed:40339580, PubMed:38600377, PubMed:38776963).

Cellular Location Cell membrane; Lipid-anchor; Cytoplasmic side

Tissue Location Expressed in fetal tissues, including testis, adrenal gland, brain, white blood cells and brain


Background

Guanine nucleotide-binding proteins (G proteins) are involved as a modulator or transducer in various transmembrane signaling systems. The beta and gamma chains are required for the GTPase activity, for replacement of GDP by GTP, and for G protein- effector interaction (By similarity).

References


Modarressi M.H.,et al.Biochem. Biophys. Res. Commun. 272:610-615(2000).
Puhl H.L. III,et al.Submitted (MAR-2002) to the EMBL/GenBank/DDBJ databases.
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Gauci S.,et al.Anal. Chem. 81:4493-4501(2009).

Images



All lanes: Anti-ROR1 antibody (C-term) at 1:250 dilution Lane 1: PANC-1 whole cell lysate Lane 2: NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 120 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

AP21903a staining GNG2 in human brain tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0. 5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6).



Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.