

MYBL2 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21915c

Product Information

Application	WB, E
Primary Accession	P10244
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB54346
Calculated MW	78764

Additional Information

Gene ID	4605
Other Names	Myb-related protein B, B-Myb, Myb-like protein 2, MYBL2, BMYB
Target/Specificity	This MYBL2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 245-278 amino acids from the Central region of human MYBL2.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MYBL2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MYBL2
Synonyms	BMYB
Function	Transcription factor involved in the regulation of cell survival, proliferation, and differentiation. Transactivates the expression of the CLU gene.
Cellular Location	Nucleus.


Background

Transcription factor involved in the regulation of cell survival, proliferation, and differentiation. Transactivates the expression of the CLU gene.

References

Nomura N.,et al.Nucleic Acids Res. 16:11075-11089(1988).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Totoki Y.,et al.Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.
Deloukas P.,et al.Nature 414:865-871(2001).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Images



All lanes : Anti-MYBL2 Antibody (Center) at 1:2000 dilution
Lane 1: Raji whole cell lysate Lane 2: HeLa whole cell lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 79 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.