

# ERVK-9 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21916b

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">Q9UKH3</a>
Other Accession	<a href="#">Q902F9</a> , <a href="#">O71037</a> , <a href="#">P61565</a> , <a href="#">P61566</a> , <a href="#">Q69384</a> , <a href="#">P61567</a> , <a href="#">Q902F8</a> , <a href="#">P63135</a>
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB54388
Calculated MW	79016

## Additional Information

---

Other Names	Endogenous retrovirus group K member 9 Env polyprotein, EnvK4 protein, Envelope polyprotein, HERV-K(C6) envelope protein, HERV-K109 envelope protein, HERV-K_6q14.1 provirus ancestral Env polyprotein, Surface protein, SU, Transmembrane protein, TM, ERVK-9
Target/Specificity	This ERVK-9 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 636-669 amino acids from human ERVK-9.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ERVK-9 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	ERVK-9
Function	Retroviral envelope proteins mediate receptor recognition and membrane fusion during early infection. Endogenous envelope proteins may have kept, lost or modified their original function during evolution. This endogenous

envelope protein has lost its original fusogenic properties.

#### Cellular Location

[Transmembrane protein]: Cell membrane; Single-pass type I membrane protein [Endogenous retrovirus group K member 9 Env polyprotein]: Virion

## Background

---

Retroviral envelope proteins mediate receptor recognition and membrane fusion during early infection. Endogenous envelope proteins may have kept, lost or modified their original function during evolution. This endogenous envelope protein has lost its original fusogenic properties.


## References

---

Barbulescu M.,et al.Curr. Biol. 9:861-868(1999).  
de Parseval N.,et al.J. Virol. 77:10414-10422(2003).  
Blaise S.,et al.Proc. Natl. Acad. Sci. U.S.A. 100:13013-13018(2003).  
Wang-Johanning F.,et al.Oncogene 22:1528-1535(2003).

## Images

---



Anti-ERVK-9 Antibody (C-Term) at dilution + ZR-75-1 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 79kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.