

TMEM115 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21934b

Product Information

Application	WB, E
Primary Accession	Q12893
Other Accession	A4FUB8 , Q9WUH1
Reactivity	Human, Mouse
Predicted	Bovine, Mouse, Rat, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB54333
Calculated MW	38197

Additional Information

Gene ID	11070
Other Names	Transmembrane protein 115, Placental protein 6, Protein PL6, TMEM115 (HGNC:30055)
Target/Specificity	This TMEM115 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 249-283 amino acids from of human TMEM115.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TMEM115 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TMEM115 (HGNC:30055)
Function	May play a role in retrograde transport of proteins from the Golgi to the endoplasmic reticulum. May indirectly play a role in protein glycosylation in the Golgi.

Cellular Location	Golgi apparatus, Golgi stack membrane; Multi-pass membrane protein
Tissue Location	Expressed strongly in kidney and skeletal muscle, followed by liver, placenta, pancreas, and lung, with low amounts in heart and only traces in brain (PubMed:11085536). Widely expressed with ubiquitous expression in epithelial tissues (at protein level) (PubMed:17973242).


Background

May play a role in retrograde transport of proteins from the Golgi to the endoplasmic reticulum. May indirectly play a role in protein glycosylation in the Golgi.

References

Lerman M.I.,et al.Cancer Res. 60:6116-6133(2000).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Muzny D.M.,et al.Nature 440:1194-1198(2006).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Ivanova A.V.,et al.J. Pathol. 214:46-57(2008).

Images



Anti-TMEM115 Antibody (C-Term) at 1:2000 dilution + human skeletal muscle lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 38 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.