

PCNXL2 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21943a

Product Information

Application WB, E **Primary Accession** A6NKB5 Reactivity Human Host Rabbit Clonality polyclonal Isotype Rabbit IgG **Clone Names** RB54379 **Calculated MW** 237277

Additional Information

Gene ID 80003

Other Names Pecanex-like protein 2, PCNXL2, KIAA0435

Target/Specificity

This PCNXL2 antibody is generated from a rabbit immunized with a KLH

conjugated synthetic peptide between 467-499 amino acids from human

PCNXL2.

Dilution WB~~1:1000-1:2000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions PCNXL2 Antibody (N-Term) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name PCNX2 (HGNC:8736)

Synonyms KIAA0435, PCNXL2

Function May play a role in tumorigenesis of colorectal carcinomas with high

microsatellite instability (MSI-H).

Cellular Location Membrane; Multi-pass membrane protein


Background

May play a role in tumorigenesis of colorectal carcinomas with high microsatellite instability (MSI-H).

References

Ishikawa K.,et al.DNA Res. 4:307-313(1997).
Ohara O.,et al.Submitted (OCT-1997) to the EMBL/GenBank/DDBJ databases.
Jikuya H.,et al.Submitted (JAN-2002) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Gregory S.G.,et al.Nature 441:315-321(2006).

Images



All lanes: Anti-PCNXL2 Antibody (N-Term) at 1:1000-1:2000 dilution Lane 1: Hela whole cell lysate Lane 2: HT-29 whole cell lysate Lane 3: WiDr whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 237 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.