

# HEXB Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21947c

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">P07686</a>
Reactivity	Human, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB54540
Calculated MW	63137

## Additional Information

---

Gene ID	3074
Other Names	Beta-hexosaminidase subunit beta, 3.2.1.52, Beta-N-acetylhexosaminidase subunit beta, Hexosaminidase subunit B, Cervical cancer proto-oncogene 7 protein, HCC-7, N-acetyl-beta-glucosaminidase subunit beta, Beta-hexosaminidase subunit beta chain B, Beta-hexosaminidase subunit beta chain A, HEXB
Target/Specificity	This HEXB antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 170-203 amino acids from the Central region of human HEXB.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	HEXB Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	HEXB ( <a href="#">HGNC:4879</a> )
Function	Hydrolyzes the non-reducing end N-acetyl-D-hexosamine and/or sulfated N-acetyl-D-hexosamine of glycoconjugates, such as the oligosaccharide

moieties from proteins and neutral glycolipids, or from certain mucopolysaccharides (PubMed:[11707436](#), PubMed:[8123671](#), PubMed:[8672428](#), PubMed:[9694901](#)). The isozyme B does not hydrolyze each of these substrates, however hydrolyzes efficiently neutral oligosaccharide (PubMed:[11707436](#)). Only the isozyme A is responsible for the degradation of GM2 gangliosides in the presence of GM2A (PubMed:[8123671](#), PubMed:[8672428](#), PubMed:[9694901](#)). During fertilization is responsible, at least in part, for the zona block to polyspermy. Present in the cortical granules of non-activated oocytes, is exocytosed during the cortical reaction in response to oocyte activation and inactivates the sperm galactosyltransferase-binding site, accounting for the block in sperm binding to the zona pellucida (By similarity).

#### Cellular Location

Lysosome. Cytoplasmic vesicle, secretory vesicle, Cortical granule {ECO:0000250|UniProtKB:P20060}


## Background

Responsible for the degradation of GM2 gangliosides, and a variety of other molecules containing terminal N-acetyl hexosamines, in the brain and other tissues.

## References

Korneluk R.G., et al. J. Biol. Chem. 261:8407-8413(1986).  
Neote K., et al. Genomics 3:279-286(1988).  
Proia R.L., et al. Proc. Natl. Acad. Sci. U.S.A. 85:1883-1887(1988).  
Kim J.W., et al. Submitted (MAY-2001) to the EMBL/GenBank/DDBJ databases.  
Kalnina N., et al. Submitted (AUG-2003) to the EMBL/GenBank/DDBJ databases.

## Images



All lanes : Anti-HEXB Antibody (Center) at 1:2000 dilution  
Lane 1: Jurkat whole cell lysate Lane 2: MCF-7 whole cell lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 63 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.