

INTS6 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21956a

Product Information

Application	WB, E
Primary Accession	Q9UL03
Other Accession	Q5JSJ4 , Q8BND4 , Q2TAF4 , Q5U4W6 , Q7SYD9 , Q6PCM2
Reactivity	Human, Rat, Mouse
Predicted	Human, Mouse, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB54384
Calculated MW	100390

Additional Information

Gene ID	26512
Other Names	Integrator complex subunit 6, Int6, DBI-1, Protein DDX26, Protein deleted in cancer 1, DICE1, INTS6, DBI1, DDX26, DDX26A
Target/Specificity	This INTS6 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 135-167 amino acids from human INTS6.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	INTS6 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	INTS6 {ECO:0000303 PubMed:33243860, ECO:0000312 HGNC:HGNC:14879}
Function	Component of the integrator complex, a multiprotein complex that terminates RNA polymerase II (Pol II) transcription in the promoter-proximal region of genes (PubMed: 33243860 , PubMed: 34004147 , PubMed: 39504960).

The integrator complex provides a quality checkpoint during transcription elongation by driving premature transcription termination of transcripts that are unfavorably configured for transcriptional elongation: the complex terminates transcription by (1) catalyzing dephosphorylation of the C-terminal domain (CTD) of Pol II subunit POLR2A and SPT5H/SPT5, (2) degrading the exiting nascent RNA transcript via endonuclease activity and (3) promoting the release of Pol II from bound DNA (PubMed:[33243860](#), PubMed:[34004147](#), PubMed:[38570683](#), PubMed:[39504960](#)). The integrator complex is also involved in terminating the synthesis of non-coding Pol II transcripts, such as enhancer RNAs (eRNAs), small nuclear RNAs (snRNAs), telomerase RNAs and long non-coding RNAs (lncRNAs) (PubMed:[16239144](#)). Within the integrator complex, INTS6 acts as a molecular adapter that promotes assembly of protein phosphatase 2A (PP2A) subunits to the integrator core complex, promoting recruitment of PP2A to transcription pause- release checkpoint (PubMed:[33243860](#), PubMed:[34004147](#)). Mediates recruitment of cytoplasmic dynein to the nuclear envelope, probably as component of the integrator complex (PubMed:[23904267](#)). May have a tumor suppressor role; an ectopic expression suppressing tumor cell growth (PubMed:[15254679](#), PubMed:[16239144](#)).

Cellular Location	Nucleus. Chromosome Note=Associates with chromatin and transcription pause-release checkpoint.
Tissue Location	Widely expressed. Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas


Background

Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes. May have a tumor suppressor role; an ectopic expression suppressing tumor cell growth.


References

Wieland I.,et al.Oncogene 18:4530-4537(1999).
 Bechtel S.,et al.BMC Genomics 8:399-399(2007).
 Dunham A.,et al.Nature 428:522-528(2004).
 Hoff H.B. III,et al.Submitted (APR-1999) to the EMBL/GenBank/DDBJ databases.
 Wieland I.,et al.Oncol. Res. 12:491-500(2001).

Images



All lanes : Anti-INTS6 Antibody (N-Term) at 1:1000 dilution
 Lane 1: HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution.
 Observed band size : 100kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-INTS6 Antibody (N-Term) at 1:2000 dilution
Lane 1: A431 whole cell lysate Lane 2: HeLa whole cell
lysate Lysates/proteins at 20 µg per lane. Secondary Goat
Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000
dilution. Predicted band size : 100 kDa Blocking/Dilution
buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.