

DNAJB4 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21964b

Product Information

Application	WB, E
Primary Accession	Q9UDY4
Other Accession	Q2KIT4 , Q5R8J8
Reactivity	Human, Mouse
Predicted	Bovine
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB54638
Calculated MW	37807

Additional Information

Gene ID	11080
Other Names	DnaJ homolog subfamily B member 4, Heat shock 40 kDa protein 1 homolog, HSP40 homolog, Heat shock protein 40 homolog, Human liver DnaJ-like protein, DNAJB4, DNAJW, HLJ1
Target/Specificity	This DNAJB4 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 216-248 amino acids from human DNAJB4.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DNAJB4 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DNAJB4
Synonyms	DNAJW, HLJ1

Function	Probable chaperone. Stimulates ATP hydrolysis and the folding of unfolded proteins mediated by HSPA1A/B (in vitro) (PubMed: 24318877).
Cellular Location	Cytoplasm. Cell membrane. Cytoplasm, myofibril, sarcomere, Z line. Note=Cytoplasmic according to PubMed:18837411 and membrane-associated according to PubMed:16542645
Tissue Location	Expressed in heart, pancreas and skeletal muscle, and to a lesser extent in brain, placenta and liver


Background

Probable chaperone.

References

Hoe K.L.,et al.Biochim. Biophys. Acta 1383:4-8(1998).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Won M.,et al.Submitted (NOV-1995) to the EMBL/GenBank/DDBJ databases.
Bienvenut W.V.,et al.Submitted (MAR-2009) to UniProtKB.

Images



Anti-DNAJB4 Antibody (C-Term) at 1:2000 dilution + HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 38 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.