

CTBS Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22021a

Product Information

Application	WB, E
Primary Accession	Q01459
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB55055
Calculated MW	43760

Additional Information

Gene ID	1486
Other Names	Di-N-acetylchitobiase, 3.2.1.-, CTBS, CTB
Target/Specificity	This CTBS antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 85-119 amino acids from of human CTBS.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CTBS Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CTBS
Synonyms	CTB
Function	Involved in the degradation of asparagine-linked glycoproteins. Hydrolyze of N-acetyl-beta-D-glucosamine (1-4)N- acetylglucosamine chitobiose core from the reducing end of the bond, it requires prior cleavage by glycosylasparaginase.

Background

Involved in the degradation of asparagine-linked glycoproteins. Hydrolyze of N-acetyl-beta-D-glucosamine (1-4)N- acetylglucosamine chitobiose core from the reducing end of the bond, it requires prior cleavage by glycosylasparaginase.

References

Fisher K.J.,et al.J. Biol. Chem. 267:19607-19616(1992).


Liu B.,et al.Glycobiology 9:589-593(1999).

Gregory S.G.,et al.Nature 441:315-321(2006).

Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Chen R.,et al.J. Proteome Res. 8:651-661(2009).

Images



Anti-CTBS Antibody (N-Term) at 1:2000 dilution + human liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 44 kDa
Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.