

# KLB Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22050a

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">Q86Z14</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB55007
Calculated MW	119808

## Additional Information

---

Gene ID	152831
Other Names	Beta-klotho, BKL, BetaKlotho, Klotho beta-like protein, KLB
Target/Specificity	This KLB antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1-29 amino acids from human KLB.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	KLB Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	KLB
Function	Contributes to the transcriptional repression of cholesterol 7-alpha-hydroxylase (CYP7A1), the rate-limiting enzyme in bile acid synthesis. Probably inactive as a glycosidase. Increases the ability of FGFR1 and FGFR4 to bind FGF21 (By similarity).
Cellular Location	Cell membrane; Single-pass type III membrane protein

## Background

---

Contributes to the transcriptional repression of cholesterol 7-alpha-hydroxylase (CYP7A1), the rate-limiting enzyme in bile acid synthesis. Probably inactive as a glycosidase. Increases the ability of FGFR1 and FGFR4 to bind FGF21 (By similarity).

## References

---


Hayashi A.,et al.Submitted (FEB-2002) to the EMBL/GenBank/DDBJ databases.

Wu X.,et al.J. Biol. Chem. 283:33304-33309(2008).

Micanovic R.,et al.J. Cell. Physiol. 219:227-234(2009).

## Images

---



All lanes : Anti-KLB Antibody (N-Term) at 1:2000 dilution  
Lane 1: A549 whole cell lysate Lane 2: HepG2 whole cell lysate Lane 3: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 120 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.