

DPYD Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22058b

Product Information

Application	WB, E
Primary Accession	Q12882
Other Accession	Q5R895
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB54986
Calculated MW	111401

Additional Information

Gene ID	1806
Other Names	Dihydropyrimidine dehydrogenase [NADP(+)], DHPDHase, DPD, 1.3.1.2, Dihydrothymine dehydrogenase, Dihydrouracil dehydrogenase, DPYD
Target/Specificity	This DPYD antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 841-875 amino acids from human DPYD.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DPYD Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DPYD (HGNC:3012)
Function	Involved in pyrimidine base degradation (PubMed: 1512248). Catalyzes the reduction of uracil and thymine (PubMed: 1512248). Also involved the degradation of the chemotherapeutic drug 5-fluorouracil (PubMed: 1512248).

Cellular Location	Cytoplasm.
Tissue Location	Found in most tissues with greatest activity found in liver and peripheral blood mononuclear cells


Background

Involved in pyrimidine base degradation. Catalyzes the reduction of uracil and thymine. Also involved the degradation of the chemotherapeutic drug 5-fluorouracil.

References

Yokota H.,et al.J. Biol. Chem. 269:23192-23196(1994).
 Johnson M.R.,et al.Cancer Res. 57:1660-1663(1997).
 Ogura K.,et al.Cancer Lett. 122:107-113(1998).
 Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
 Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



Anti-DPYD Antibody (C-Term) at 1:2000 dilution + mouse lung lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 111 kDa
 Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.