

# RPL13 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22061c

## Product Information

---

|                          |   |
|--------------------------|---|
| <b>Application</b>       | WB, E   |
| <b>Primary Accession</b> | <a href="#">P26373</a>  |
| <b>Other Accession</b>   | <a href="#">Q56JZ1</a> , <a href="#">Q9Z313</a> , <a href="#">P47963</a> , <a href="#">P41123</a> |
| <b>Reactivity</b>        | Human, Rat, Mouse   |
| <b>Predicted</b>         | Bovine, Mouse, Rat  |
| <b>Host</b>              | Rabbit  |
| <b>Clonality</b>         | polyclonal  |
| <b>Isotype</b>           | Rabbit IgG  |
| <b>Clone Names</b>       | RB55057   |
| <b>Calculated MW</b>     | 24261   |

## Additional Information

---

|                           |  |
|---------------------------|--|
| <b>Gene ID</b>            | 6137   |
| <b>Other Names</b>        | 60S ribosomal protein L13, Breast basic conserved protein 1, RPL13, BBC1   |
| <b>Target/Specificity</b> | This RPL13 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 49-82 amino acids from the Central region of human RPL13.           |
| <b>Dilution</b>           | WB~~1:2000 E~~Use at an assay dependent concentration.   |
| <b>Format</b>             | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| <b>Storage</b>            | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.                                      |
| <b>Precautions</b>        | RPL13 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.  |

## Protein Information

---

|                 |  |
|-----------------|--|
| <b>Name</b>     | RPL13  |
| <b>Synonyms</b> | BBC1   |
| <b>Function</b> | Component of the ribosome, a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell (PubMed: <a href="#">23636399</a> , |

PubMed:[31630789](#), PubMed:[32669547](#)). The small ribosomal subunit (SSU) binds messenger RNAs (mRNAs) and translates the encoded message by selecting cognate aminoacyl-transfer RNA (tRNA) molecules (Probable). The large subunit (LSU) contains the ribosomal catalytic site termed the peptidyl transferase center (PTC), which catalyzes the formation of peptide bonds, thereby polymerizing the amino acids delivered by tRNAs into a polypeptide chain (Probable). The nascent polypeptides leave the ribosome through a tunnel in the LSU and interact with protein factors that function in enzymatic processing, targeting, and the membrane insertion of nascent chains at the exit of the ribosomal tunnel (Probable). As part of the LSU, it is probably required for its formation and the maturation of rRNAs (PubMed:[31630789](#)). Plays a role in bone development (PubMed:[31630789](#)).


**Cellular Location** Cytoplasm

**Tissue Location** Higher levels of expression in benign breast lesions than in carcinomas.

## References

Adams S.M.,et al.Hum. Mol. Genet. 1:91-96(1992).  
 Shichijo S.,et al.Submitted (MAY-2001) to the EMBL/GenBank/DDBJ databases.  
 Ota T.,et al.Nat. Genet. 36:40-45(2004).  
 Martin J.,et al.Nature 432:988-994(2004).  
 Daub H.,et al.Mol. Cell 31:438-448(2008).

## Images



All lanes : Anti-RPL13 Antibody (Center) at 1:2000 dilution  
 Lane 1: 293T/17 whole cell lysate Lane 2: HeLa whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: K562 whole cell lysate Lane 5: MCF-7 whole cell lysate  
 Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.