

# TFF1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP22076c

## Product Information

---

<b>Application</b>	WB, IF, E
<b>Primary Accession</b>	<a href="#">P04155</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB55405
<b>Calculated MW</b>	9150

## Additional Information

---

<b>Gene ID</b>	7031
<b>Other Names</b>	Trefoil factor 1, Breast cancer estrogen-inducible protein, PNR-2, Polypeptide P1.A, hP1.A, Protein pS2, TFF1, BCEI, PS2
<b>Target/Specificity</b>	This TFF1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 11-40 amino acids from the Central region of human TFF1.
<b>Dilution</b>	WB~~1:2000 IF~~1:25 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	TFF1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	TFF1
<b>Synonyms</b>	BCEI, PS2
<b>Function</b>	Stabilizer of the mucous gel overlying the gastrointestinal mucosa that provides a physical barrier against various noxious agents. May inhibit the growth of calcium oxalate crystals in urine.

**Cellular Location**

Secreted

**Tissue Location**


Found in stomach, with highest levels in the upper gastric mucosal cells (at protein level). Detected in goblet cells of the small and large intestine and rectum, small submucosal glands in the esophagus, mucous acini of the sublingual gland, submucosal glands of the trachea, and epithelial cells lining the exocrine pancreatic ducts but not in the remainder of the pancreas (at protein level) Scattered expression is detected in the epithelial cells of the gallbladder and submucosal glands of the vagina, and weak expression is observed in the bronchial goblet cells of the pseudostratified epithelia in the respiratory system (at protein level). Detected in urine (at protein level). Strongly expressed in breast cancer but at low levels in normal mammary tissue. It is regulated by estrogen in MCF-7 cells. Strong expression found in normal gastric mucosa and in the regenerative tissues surrounding ulcerous lesions of gastrointestinal tract, but lower expression found in gastric cancer (at protein level).

**Background**

Stabilizer of the mucous gel overlying the gastrointestinal mucosa that provides a physical barrier against various noxious agents. May inhibit the growth of calcium oxalate crystals in urine.


**References**

- Jakowlew S.B., et al. *Nucleic Acids Res.* 12:2861-2878(1984).  
Prud'Homme J.-F., et al. *DNA* 4:11-21(1985).  
Jeltsch J.-M., et al. *Nucleic Acids Res.* 15:1401-1414(1987).  
Takahashi H., et al. *FEBS Lett.* 261:283-286(1990).  
Mori K., et al. *J. Biochem.* 107:73-76(1990).

**Images**

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized MCF-7 (human breast cancer cell line) cells labeling TFF1 with AP22076c at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG (NK179883) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing secreted staining on MCF-7 cell line. The nuclear counter stain is DAPI (blue).

All lanes : Anti-TFF1 Antibody (Center) at 1:2000 dilution  
Lane 1: Jurkat whole cell lysate Lane 2: T47D whole cell lysate Lane 3: SK-BR-3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 9 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.