10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

C8orf80 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22085a

Product Information

Application	WB, E
Primary Accession	Q68CJ6
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB55346
Calculated MW	91132

Additional Information

Gene ID	389643
Other Names	Nuclear GTPase SLIP-GC, 3.6.1.-, Speckled-like pattern in the germinal center, NUGGC, C8orf80
Target/Specificity	This C8orf80 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 27-59 amino acids from human C8orf80.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	C8orf80 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NUGGC
Synonyms	C8orf80
Function	Nuclear GTPase found in germinal center B-cells, where it may inhibit function of the activation-induced cytidine deaminase AICDA

	(PubMed:19734146). Reduces somatic hypermutation in B-cells which may enhance genome stability (By similarity).
Cellular Location	Nucleus speckle.
Tissue Location	Expressed in germinal center B-cell and in lymphomas derived from germinal center B-cell


Background

Plays a role as replication-related GTPase protein in germinal center B-cell.

References

Nusbaum C.,et al.Nature 439:331-335(2006).
Yamada S.,et al.Oncogene 23:5901-5911(2004).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Richter K.,et al.J. Biol. Chem. 284:30652-30661(2009).

Images



Anti-C8orf80 Antibody (N-Term) at 1:2000 dilution + Ramos whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 91 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.