10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

PIGC Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22091b

Product Information

Application	WB, FC, E
Primary Accession	Q92535
Other Accession	Q3ZBX1, Q9CXR4, Q5PQQ4
Reactivity	Human, Rat, Mouse
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB54704
Calculated MW	33583

Additional Information

Gene ID	5279
Other Names	Phosphatidylinositol N-acetylglucosaminyltransferase subunit C, 2.4.1.198, Phosphatidylinositol-glycan biosynthesis class C protein, PIG-C, PIGC, GPI2
Target/Specificity	This PIGC antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 264-294 amino acids from human PIGC.
Dilution	WB~~1:1000 FC~~1:25 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PIGC Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PIGC (HGNC:8960)
Synonyms	GPI2
Function	Part of the glycosylphosphatidylinositol-N- acetylglucosaminyltransferase

(GPI-GnT) complex that catalyzes the transfer of N-acetylglucosamine from UDP-N-acetylglucosamine to phosphatidylinositol and participates in the first step of GPI biosynthesis.

Cellular Location Endoplasmic reticulum membrane; Multi-pass membrane protein


Background

Part of the complex catalyzing the transfer of N- acetylglucosamine from UDP-N-acetylglucosamine to phosphatidylinositol, the first step of GPI biosynthesis.


References

Inoue N.,et al.Biochem. Biophys. Res. Commun. 226:193-199(1996).
Hong Y.,et al.Genomics 44:347-349(1997).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Ebert L.,et al.Submitted (MAY-2004) to the EMBL/GenBank/DDBJ databases.
Gregory S.G.,et al.Nature 441:315-321(2006).

Images



Overlay histogram showing K562 cells stained with AP22091b (green line). The cells were fixed with 2% paraformaldehyde (10 min). The cells were then icubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP22091b, 1:25 dilution) for 60 min at 37ºC. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OH191631) at 1/200 dilution for 40 min at 37ºC. Isotype control antibody (blue line) was rabbit IgG (1µg/1x10^6 cells) used under the same conditions. Acquisition of >10, 000 events was performed.



All lanes : Anti-PIGC Antibody (C-Term) at 1:1000 dilution Lane 1: K562 whole cell lysate Lane 2: MOLT-4 whole cell lysate Lane 3: PC-3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 34 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.