10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ALG6 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22095c

Product Information

Application	WB, E
Primary Accession	Q9Y672
Other Accession	Q3TAE8, Q5NVS8, Q3T1L5
Reactivity	Human, Rat, Mouse
Predicted	Mouse, Rat, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB55877
Calculated MW	58121

Additional Information

Gene ID	29929
Other Names	Dolichyl pyrophosphate Man9GlcNAc2 alpha-1, 3-glucosyltransferase, 2.4.1.267, Asparagine-linked glycosylation protein 6 homolog, Dol-P-Glc:Man(9)GlcNAc(2)-PP-Dol alpha-1, 3-glucosyltransferase, Dolichyl-P-Glc:Man9GlcNAc2-PP-dolichyl glucosyltransferase, ALG6
Target/Specificity	This ALG6 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 241-274 amino acids from the Central region of human ALG6.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ALG6 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ALG6 (HGNC:23157)
Function	Dolichyl pyrophosphate Man9GlcNAc2 alpha-1,3- glucosyltransferase that

operates in the biosynthetic pathway of dolichol-linked oligosaccharides, the glycan precursors employed in protein asparagine (N)-glycosylation. The assembly of dolichol-linked oligosaccharides begins on the cytosolic side of the endoplasmic reticulum membrane and finishes in its lumen. The sequential addition of sugars to dolichol pyrophosphate produces dolichol-linked oligosaccharides containing fourteen sugars, including two GlcNAcs, nine mannoses and three glucoses. Once assembled, the oligosaccharide is transferred from the lipid to nascent proteins by oligosaccharyltransferases. In the lumen of the endoplasmic reticulum, adds the first glucose residue from dolichyl phosphate glucose (Dol-P- Glc) onto the lipid-linked oligosaccharide intermediate Man(9)GlcNAc(2)-PP-Dol to produce Glc(1)Man(9)GlcNAc(2)-PP-Dol. Glc(1)Man(9)GlcNAc(2)-PP-Dol is a substrate for ALG8, the following enzyme in the biosynthetic pathway.

Cellular Location Endoplasmic reticulum membrane; Multi-pass membrane protein


Background

Adds the first glucose residue to the lipid-linked oligosaccharide precursor for N-linked glycosylation. Transfers glucose from dolichyl phosphate glucose (Dol-P-Glc) onto the lipid-linked oligosaccharide Man(9)GlcNAc(2)-PP-Dol.

References

Imbach T.,et al.Proc. Natl. Acad. Sci. U.S.A. 96:6982-6987(1999).
Mao Y.M.,et al.Submitted (MAY-1998) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Gregory S.G.,et al.Nature 441:315-321(2006).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Images



All lanes : Anti-ALG6 Antibody (Center) at 1:2000 dilution Lane 1: human fetal lung lysate Lane 2: human fetal heart lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 58 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.