

CSRNP1 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22135b

Product Information

Application	WB, E
Primary Accession	Q96S65
Other Accession	Q5R638
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB55909
Calculated MW	63508

Additional Information

Gene ID	64651
Other Names	Cysteine/serine-rich nuclear protein 1, CSRNP-1, Axin-1 up-regulated gene 1 protein, Protein URAX1, TGF-beta-induced apoptosis protein 3, TAIP-3, CSRNP1, AXUD1, TAIP3
Target/Specificity	This CSRNP1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 501-533 amino acids from human CSRNP1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CSRNP1 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CSRNP1
Synonyms	AXUD1, TAIP3
Function	Binds to the consensus sequence 5'-AGAGTG-3' and has transcriptional

activator activity (By similarity). May have a tumor-suppressor function. May play a role in apoptosis.

Cellular Location

Nucleus.

Tissue Location

Ubiquitous. Most abundantly expressed in lung, placenta, skeletal muscle, pancreas and leukocyte. Frequently down-regulated in lung, kidney, liver and colon cancers compared with their corresponding normal tissues.


Background

Binds to the consensus sequence 5'-AGAGTG-3' and has transcriptional activator activity (By similarity). May have a tumor-suppressor function. May play a role in apoptosis.

References

Ishiguro H., et al. Oncogene 20:5062-5066(2001).
Akiyama N., et al. Submitted (JUN-2001) to the EMBL/GenBank/DDBJ databases.
Muzny D.M., et al. Nature 440:1194-1198(2006).
Bechtel S., et al. BMC Genomics 8:399-399(2007).

Images



Anti-CSRNP1 Antibody (C-Term) at 1:1000 dilution + human lung whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 64 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.