

SLC2A3 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP22174b

Product Information

Application WB, FC, E **Primary Accession** P11169

Other Accession Q8TDB8, Q5R608, Q9XSC2
Reactivity Human, Mouse, Rat

Predicted Human
Host Rabbit
Clonality polyclonal
Isotype Rabbit IgG
Clone Names RB56266
Calculated MW 53924

Additional Information

Gene ID 6515

Other Names Solute carrier family 2, facilitated glucose transporter member 3, Glucose

transporter type 3, brain, GLUT-3, SLC2A3, GLUT3

Target/Specificity This SLC2A3 antibody is generated from a rabbit immunized with a KLH

conjugated synthetic peptide between 432-463 amino acids from human

SLC2A3.

Dilution WB~~1:2000 FC~~1:25 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This

antibody is purified through a protein A column, followed by peptide affinity

purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions SLC2A3 Antibody (C-Term) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name SLC2A3 (HGNC:11007)

Function Facilitative glucose transporter (PubMed: <u>26176916</u>, PubMed: <u>32860739</u>,

PubMed:<u>9477959</u>). Can also mediate the uptake of various other monosaccharides across the cell membrane (PubMed:<u>26176916</u>,

PubMed:<u>9477959</u>). Mediates the uptake of glucose, 2- deoxyglucose, galactose, mannose, xylose and fucose, and probably also dehydroascorbate (PubMed:<u>26176916</u>, PubMed:<u>9477959</u>). Does not mediate fructose transport (PubMed:<u>26176916</u>, PubMed:<u>9477959</u>). Required for mesendoderm

differentiation (By similarity).

Cellular Location Cell membrane; Multi-pass membrane protein. Perikaryon

{ECO:0000250 | UniProtKB:Q07647}. Cell projection

{ECO:0000250|UniProtKB:Q07647}. Note=Localized to densely spaced patches

along neuronal processes. {ECO:0000250 | UniProtKB:Q07647}

Tissue Location Highly expressed in brain (PubMed:8457197). Expressed in many tissues.

Background

Facilitative glucose transporter. Probably a neuronal glucose transporter.

References

Kayano T.,et al.J. Biol. Chem. 263:15245-15248(1988). Stuart C.A.,et al.Submitted (JUN-2000) to the EMBL/GenBank/DDBJ databases. Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases. Ota T.,et al.Nat. Genet. 36:40-45(2004). Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.