

EPHX2 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP22178a

Product Information

Primary AccessionP34913ReactivityHumanHostRabbitClonalitypolyclonalIsotypeRabbit IgGClone NamesRB56273Calculated MW62616	Application	WB, E
ReactivityHumanHostRabbitClonalitypolyclonalIsotypeRabbit IgGClone NamesRB56273Calculated MW62616	Primary Accession	<u>P34913</u>
HostRabbitClonalitypolyclonalIsotypeRabbit IgGClone NamesRB56273Calculated MW62616	Reactivity	Human
ClonalitypolyclonalIsotypeRabbit IgGClone NamesRB56273Calculated MW62616	Host	Rabbit
IsotypeRabbit IgGClone NamesRB56273Calculated MW62616	Clonality	polyclonal
Clone NamesRB56273Calculated MW62616	Isotype	Rabbit IgG
Calculated MW 62616	Clone Names	RB56273
	Calculated MW	62616

Additional Information

Gene ID	2053
Other Names	Bifunctional epoxide hydrolase 2, Cytosolic epoxide hydrolase 2, CEH, 3.3.2.10, Epoxide hydratase, Soluble epoxide hydrolase, SEH, Lipid-phosphate phosphatase, 3.1.3.76, EPHX2
Target/Specificity	This EPHX2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 64-94 amino acids from human EPHX2.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	EPHX2 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	EPHX2 (<u>HGNC:3402</u>)
Function	Bifunctional enzyme (PubMed: <u>12574510</u>). The C-terminal domain has epoxide hydrolase activity and acts on epoxides (alkene oxides, oxiranes) and arene oxides (PubMed: <u>12574510</u> , PubMed: <u>12869654</u> , PubMed: <u>22798687</u>). Plays a role in xenobiotic metabolism by degrading potentially toxic epoxides (By similarity). Also determines steady- state levels of physiological mediators

(PubMed:<u>12574510</u>, PubMed:<u>12869654</u>, PubMed:<u>21217101</u>, PubMed:<u>22798687</u>).

Cellular Location

Cytoplasm. Peroxisome.

Background

Bifunctional enzyme. The C-terminal domain has epoxide hydrolase activity and acts on epoxides (alkene oxides, oxiranes) and arene oxides. Plays a role in xenobiotic metabolism by degrading potentially toxic epoxides. Also determines steady-state levels of physiological mediators. The N-terminal domain has lipid phosphatase activity, with the highest activity towards threo- 9,10-phosphonooxy-hydroxy-octadecanoic acid, followed by erythro- 9,10-phosphonooxy-hydroxy-octadecanoic acid, 12-phosphonooxy-octadec-9E-enoic acid, and p-nitrophenyl phospate.

References

Beetham J.K.,et al.Arch. Biochem. Biophys. 305:197-201(1993). Sandberg M.,et al.Biochem. Biophys. Res. Commun. 221:333-339(1996). Sandberg M.,et al.J. Biol. Chem. 275:28873-28881(2000). Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases. Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



All lanes : Anti-EPHX2 Antibody (N-Term) at 1:2000 dilution Lane 1: Jurkat whole cell lysate Lane 2: human fetal kidney lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 63 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.