

KIAA1524 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22183a

Product Information

Application	WB, E
Primary Accession	Q8TCG1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB56267
Calculated MW	102185

Additional Information

Gene ID	57650
Other Names	Protein CIP2A, Cancerous inhibitor of PP2A, p90 autoantigen, KIAA1524, CIP2A
Target/Specificity	This KIAA1524 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 269-301 amino acids from human KIAA1524.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	KIAA1524 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CIP2A {ECO:0000303 PubMed:17632056, ECO:0000312 HGNC:HGNC:29302}
Function	Acts as an inhibitor of protein phosphatase PP2A (PubMed: 17632056). Promotes anchorage-independent cell growth and tumor formation by preventing dephosphorylation of MYC, thereby stabilizing MYC in human malignancies (PubMed: 17632056). Together with TOPBP1, plays an essential role in the response to genome instability generated by the presence of

acentric chromosome fragments derived from shattered chromosomes within micronuclei (PubMed:[35121901](#), PubMed:[35842428](#), PubMed:[37165191](#), PubMed:[37316668](#)). Micronuclei, which are frequently found in cancer cells, consist of chromatin surrounded by their own nuclear membrane: following breakdown of the micronuclear envelope, a process associated with chromothripsis, the CIP2A-TOPBP1 complex tethers chromosome fragments during mitosis to ensure clustered segregation of the fragments to a single daughter cell nucleus, facilitating re-ligation with limited chromosome scattering and loss (PubMed:[37165191](#), PubMed:[37316668](#)).

Cellular Location

Cytoplasm. Chromosome. Note=Predominantly localizes within the cytoplasm (PubMed:[35842428](#)). Localizes to broken chromosomes within micronuclei during interphase and following chromothripsis (PubMed:[37165191](#), PubMed:[37316668](#)). Localization to broken chromosomes is mainly independent of MDC1 (PubMed:[35121901](#), PubMed:[37165191](#))

Tissue Location

Expressed at low levels in most of the tissues. Overexpressed in head-and-neck squamous cell carcinomas (HNSCC) Present in liver cancer cells (at protein level)


Background

Oncoprotein that inhibits PP2A and stabilizes MYC in human malignancies. Promotes anchorage-independent cell growth and tumor formation.

References

Soo Hoo L.,et al.Oncogene 21:5006-5015(2002).
Nagase T.,et al.DNA Res. 7:143-150(2000).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Shi F.D.,et al.Prostate 63:252-258(2005).

Images



Anti-KIAA1524 Antibody (N-Term) at 1:2000 dilution + mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 102 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.