10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

UHRF1BP1 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22184b

Product Information

Application	WB, E
Primary Accession	Q6BDS2
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB56270
Calculated MW	159485

Additional Information

Gene ID	54887
Other Names	UHRF1-binding protein 1, ICBP90-binding protein 1, Ubiquitin-like containing PHD and RING finger domains 1-binding protein 1, UHRF1BP1, C6orf107
Target/Specificity	This UHRF1BP1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1096-1129 amino acids from the human UHRF1BP1.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	UHRF1BP1 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	BLTP3A (HGNC:21216)
Synonyms	C6orf107, UHRF1BP1
Function	Tube-forming lipid transport protein which probably mediates the transfer of lipids between membranes at organelle contact sites (PubMed:35499567).

May be involved in the retrograde traffic of vesicle clusters in the endocytic pathway to the Golgi complex (PubMed:35499567).

Cellular Location Late endosome.


Background

May act as a negative regulator of cell growth.

References

Unoki M.,et al.Oncogene 23:7601-7610(2004).
Mungall A.J.,et al.Nature 425:805-811(2003).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).
Gauci S.,et al.Anal. Chem. 81:4493-4501(2009).

Images



Anti-UHRF1BP1 Antibody (C-Term) at 1:2000 dilution + A431 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 159 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.