

# UHRF1BP1 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22184b

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">Q6BDS2</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB56270
<b>Calculated MW</b>	159485

## Additional Information

---

<b>Gene ID</b>	54887
<b>Other Names</b>	UHRF1-binding protein 1, ICBP90-binding protein 1, Ubiquitin-like containing PHD and RING finger domains 1-binding protein 1, UHRF1BP1, C6orf107
<b>Target/Specificity</b>	This UHRF1BP1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1096-1129 amino acids from the human UHRF1BP1.
<b>Dilution</b>	WB~~1:2000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	UHRF1BP1 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	BLTP3A ( <a href="#">HGNC:21216</a> )
<b>Synonyms</b>	C6orf107, UHRF1BP1
<b>Function</b>	Tube-forming lipid transport protein which probably mediates the transfer of lipids between membranes at organelle contact sites (PubMed: <a href="#">35499567</a> ). May be involved in the retrograde traffic of vesicle clusters in the endocytic

pathway to the Golgi complex (PubMed:[35499567](#)).

## Cellular Location

Late endosome.


## Background

May act as a negative regulator of cell growth.

## References

Unoki M.,et al.Oncogene 23:7601-7610(2004).  
Mungall A.J.,et al.Nature 425:805-811(2003).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).  
Gauci S.,et al.Anal. Chem. 81:4493-4501(2009).

## Images



Anti-UHRF1BP1 Antibody (C-Term) at 1:2000 dilution +  
A431 whole cell lysate Lysates/proteins at 20  $\mu$ g per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase  
conjugated at 1/10000 dilution. Predicted band size : 159  
kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.