

# TOMM40 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22253a

## Product Information

---

<b>Application</b>	WB, FC, IF, E
<b>Primary Accession</b>	<a href="#">O96008</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB56900
<b>Calculated MW</b>	37893

## Additional Information

---

<b>Gene ID</b>	10452
<b>Other Names</b>	Mitochondrial import receptor subunit TOM40 homolog, Protein Haymaker, Translocase of outer membrane 40 kDa subunit homolog, p38.5, TOMM40, C19orf1, PEREC1, TOM40
<b>Target/Specificity</b>	This TOMM40 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 22-56 amino acids from human TOMM40.
<b>Dilution</b>	WB~~1:2000 FC~~1:25 IF~~1:25 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	TOMM40 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	TOMM40
<b>Synonyms</b>	C19orf1, PEREC1, TOM40
<b>Function</b>	Channel-forming protein essential for import of protein precursors into mitochondria (PubMed: <a href="#">15644312</a> , PubMed: <a href="#">31206022</a> ). Plays a role in the

assembly of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) by forming a complex with BCAP31 and mediating the translocation of Complex I components from the cytosol to the mitochondria (PubMed:[31206022](#)).

## Cellular Location

Mitochondrion outer membrane; Multi-pass membrane protein.  
Note=Associates with the mitochondria- associated ER membrane via interaction with BCAP31


## Background

Channel-forming protein essential for import of protein precursors into mitochondria.


## References

Freitas E.M.,et al.DNA Seq. 9:89-100(1998).  
Yoshiura K.,et al.Submitted (JAN-1998) to the EMBL/GenBank/DDBJ databases.  
Das B.,et al.Int. J. Cancer 94:800-806(2001).  
Lubec G.,et al.Submitted (MAR-2007) to UniProtKB.  
Johnston A.J.,et al.J. Biol. Chem. 277:42197-42204(2002).

## Images




Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized U-2 OS (human osteosarcoma cell line) cells labeling TOMM40 with AP22253a at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG (1583138) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing mitochondrion staining on U-2 OS cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin (PD18466410) at 1/100 dilution (red).The nuclear counter stain is DAPI (blue).



Overlay histogram showing HeLa cells stained with AP22253a(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP22253a, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(1583138) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10, 000 events was performed.

All lanes : Anti-TOMM40 Antibody (N-Term) at 1:2000 dilution  
Lane 1: Human brain lysate  
Lane 2: 293 whole cell lysate  
Lane 3: U-87 MG whole cell lysate  
Lane 4: Hut78 whole cell lysate  
Lane 5: Hela whole cell lysate  
Lane 6: A431 whole cell lysate  
Lane 7: HepG2 whole cell



lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 38 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.