

SSH3 (Ser37) Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22265a

Product Information

Application	WB, E
Primary Accession	Q8TE77
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB56878
Calculated MW	72996

Additional Information

Gene ID	54961
Other Names	Protein phosphatase Slingshot homolog 3, 3.1.3.16, 3.1.3.48, SSH-like protein 3, SSH-3L, hSSH-3L, SSH3, SSH3L
Target/Specificity	This SSH3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 10-43 amino acids from human SSH3.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SSH3 (Ser37) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SSH3
Synonyms	SSH3L
Function	Protein phosphatase which may play a role in the regulation of actin filament dynamics. Can dephosphorylate and activate the actin binding/depolymerizing factor cofilin, which subsequently binds to actin filaments and stimulates their disassembly (By similarity).


Background

Protein phosphatase which may play a role in the regulation of actin filament dynamics. Can dephosphorylate and activate the actin binding/depolymerizing factor cofilin, which subsequently binds to actin filaments and stimulates their disassembly (By similarity).

References

Niwa R.,et al.Cell 108:233-246(2002).
Ohta Y.,et al.Genes Cells 8:811-824(2003).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Matsuoka S.,et al.Science 316:1160-1166(2007).
Daub H.,et al.Mol. Cell 31:438-448(2008).

Images



Anti-SSH3 (Ser37) Antibody at 1:2000 dilution + ZR-75-1 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 73 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.