

# MYPT1 (Ser668) Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP22268a

## **Product Information**

Application WB, E Primary Accession 014974

Other Accession

Reactivity

Predicted

Q90623, Q9DBR7, Q10728

Human, Mouse, Rat

Mouse, Rat, Chicken

Host Rabbit
Clonality polyclonal
Isotype Rabbit IgG
Clone Names RB56874
Calculated MW 115281

#### **Additional Information**

**Gene ID** 4659

Other Names Protein phosphatase 1 regulatory subunit 12A, Myosin phosphatase-targeting

subunit 1, Myosin phosphatase target subunit 1, Protein phosphatase

myosin-binding subunit, PPP1R12A, MBS, MYPT1

**Target/Specificity** This MYPT1 (Ser668) antibody is generated from a rabbit immunized with a

KLH conjugated synthetic peptide between 641-674 amino acids from human

MYPT1.

**Dilution** WB~~1:4000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** MYPT1 (Ser668) Antibody is for research use only and not for use in diagnostic

or therapeutic procedures.

#### **Protein Information**

Name PPP1R12A ( <u>HGNC:7618</u>)

**Function** Key regulator of protein phosphatase 1C (PPP1C). Mediates binding to

myosin. As part of the PPP1C complex, involved in dephosphorylation of

PLK1. Capable of inhibiting HIF1AN-dependent suppression of HIF1A activity.

**Cellular Location** Cytoplasm, cytoskeleton, stress fiber. Note=Also along actomyosin

filaments

**Tissue Location** Expressed in striated muscles, specifically in type 2a fibers (at protein level).


# **Background**

Key regulator of protein phosphatase 1C (PPP1C). Mediates binding to myosin. As part of the PPP1C complex, involved in dephosphorylation of PLK1. Capable of inhibiting HIF1AN- dependent suppression of HIF1A activity.

#### References

Takahashi N.,et al.Genomics 44:150-152(1997). Guo J.H.,et al.Submitted (DEC-2001) to the EMBL/GenBank/DDBJ databases. Xia D.,et al.Submitted (SEP-2003) to the EMBL/GenBank/DDBJ databases. Ota T.,et al.Nat. Genet. 36:40-45(2004). Scherer S.E.,et al.Nature 440:346-351(2006).

### **Images**



All lanes: Anti-MYPT1 (Ser668) Antibody at 1:4000 dilution Lane 1: Hela whole cell lysate Lane 2: MCF-7 whole cell lysate Lane 3: C2C12 whole cell lysate Lane 4: NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 115 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.