

MST1R Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22294c

Product Information

Application	WB, E
Primary Accession	Q04912
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB58004
Calculated MW	152241

Additional Information

Gene ID	4486
Other Names	Macrophage-stimulating protein receptor, MSP receptor, 2.7.10.1, CDw136, Protein-tyrosine kinase 8, p185-Ron, CD136, Macrophage-stimulating protein receptor alpha chain, Macrophage-stimulating protein receptor beta chain, MST1R, PTK8, RON
Target/Specificity	This MST1R antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 600-633 amino acids from the Central region of human MST1R.
Dilution	WB~~1:1000-1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MST1R Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MST1R
Synonyms	PTK8, RON

Function	Receptor tyrosine kinase that transduces signals from the extracellular matrix into the cytoplasm by binding to MST1 ligand. Regulates many physiological processes including cell survival, migration and differentiation. Ligand binding at the cell surface induces autophosphorylation of RON on its intracellular domain that provides docking sites for downstream signaling molecules. Following activation by ligand, interacts with the PI3-kinase subunit PIK3R1, PLCG1 or the adapter GAB1. Recruitment of these downstream effectors by RON leads to the activation of several signaling cascades including the RAS-ERK, PI3 kinase-AKT, or PLCgamma-PKC. RON signaling activates the wound healing response by promoting epithelial cell migration, proliferation as well as survival at the wound site. Also plays a role in the innate immune response by regulating the migration and phagocytic activity of macrophages. Alternatively, RON can also promote signals such as cell migration and proliferation in response to growth factors other than MST1 ligand.
Cellular Location	Membrane; Single-pass type I membrane protein.
Tissue Location	Expressed in colon, skin, lung and bone marrow.


Background

Receptor tyrosine kinase that transduces signals from the extracellular matrix into the cytoplasm by binding to MST1 ligand. Regulates many physiological processes including cell survival, migration and differentiation. Ligand binding at the cell surface induces autophosphorylation of RON on its intracellular domain that provides docking sites for downstream signaling molecules. Following activation by ligand, interacts with the PI3-kinase subunit PIK3R1, PLCG1 or the adapter GAB1. Recruitment of these downstream effectors by RON leads to the activation of several signaling cascades including the RAS-ERK, PI3 kinase-AKT, or PLCgamma-PKC. RON signaling activates the wound healing response by promoting epithelial cell migration, proliferation as well as survival at the wound site. Plays also a role in the innate immune response by regulating the migration and phagocytic activity of macrophages. Alternatively, RON can also promote signals such as cell migration and proliferation in response to growth factors other than MST1 ligand.

References

Ronsin C.,et al.Oncogene 8:1195-1202(1993).
Collesi C.,et al.Mol. Cell. Biol. 16:5518-5526(1996).
Jin P.,et al.Arthritis Res. Ther. 10:R73-R73(2008).
Muzny D.M.,et al.Nature 440:1194-1198(2006).
Ponzetto C.,et al.Mol. Cell. Biol. 13:4600-4608(1993).

Images



All lanes : Anti-MST1R Antibody (Center) at 1:1000-1:2000 dilution
Lane 1: HACAT whole cell lysate
Lane 2: HCT116 whole cell lysate
Lane 3: SK-BR-3 whole cell lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 152 kDa
Blocking/Dilution buffer: 5% NFDN/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.