

DEPTOR Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22297a

Product Information

Application	WB, E
Primary Accession	Q8TB45
Other Accession	Q570Y9
Reactivity	Human, Mouse
Predicted	Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB57442
Calculated MW	46294

Additional Information

Gene ID	64798
Other Names	DEP domain-containing mTOR-interacting protein, DEP domain-containing protein 6, DEPTOR, DEPDC6
Target/Specificity	This DEPTOR antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 86-120 amino acids from the human region of human DEPTOR.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DEPTOR Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DEPTOR {ECO:0000303 PubMed:19446321, ECO:0000312 HGNC:HGNC:22953}
Function	Negative regulator of the mTORC1 and mTORC2 complexes: inhibits the protein kinase activity of MTOR, thereby inactivating both complexes

(PubMed:[19446321](#), PubMed:[22017875](#), PubMed:[22017876](#), PubMed:[22017877](#), PubMed:[25936805](#), PubMed:[29382726](#), PubMed:[34519268](#), PubMed:[34519269](#)). DEPTOR inhibits mTORC1 and mTORC2 to induce autophagy (PubMed:[22017875](#), PubMed:[22017876](#), PubMed:[22017877](#)). In contrast to AKT1S1/PRAS40, only partially inhibits mTORC1 activity (PubMed:[34519268](#), PubMed:[34519269](#)).

Cellular Location

Lysosome membrane. Note=Localizes to the lysosomal membrane when associated with the mTORC1 and mTORC2 complexes.


Background

Negative regulator of the mTORC1 and mTORC2 signaling pathways. Inhibits the kinase activity of both complexes.

References

Wiemann S.,et al.Genome Res. 11:422-435(2001).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Nusbaum C.,et al.Nature 439:331-335(2006).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Gauci S.,et al.Anal. Chem. 81:4493-4501(2009).

Images



Anti-DEPTOR Antibody (N-Term) at 1:2000 dilution + 293 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 46 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.