10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MUT Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22355c

Product Information

Application	WB, IF, E
Primary Accession	P22033
Other Accession	Q5RFN2
Reactivity	Human, Mouse
Predicted	Human, Rat, Bovine, Rabbit
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB57881
Calculated MW	83134

Additional Information

Gene ID	4594
Other Names	Methylmalonyl-CoA mutase, mitochondrial, MCM, 5.4.99.2, Methylmalonyl-CoA isomerase, MUT
Target/Specificity	This MUT antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 492-526 amino acids from the Central region of human MUT.
Dilution	WB~~1:2000 IF~~1:25 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MUT Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MMUT (HGNC:7526)
Function	Catalyzes the reversible isomerization of methylmalonyl-CoA (MMCoA) (generated from branched-chain amino acid metabolism and degradation of dietary odd chain fatty acids and cholesterol) to succinyl-CoA

(3-carboxypropionyl-CoA), a key intermediate of the tricarboxylic acid cycle.

Cellular Location Mitochondrion matrix. Mitochondrion. Cytoplasm


Background

Involved in the degradation of several amino acids, odd- chain fatty acids and cholesterol via propionyl-CoA to the tricarboxylic acid cycle. MCM has different functions in other species.


References

Jansen R.,et al.Genomics 4:198-205(1989).
Nham S.U.,et al.Genomics 8:710-716(1990).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Mungall A.J.,et al.Nature 425:805-811(2003).

Images




Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized Hela (human cervical epithelial adenocarcinoma cell line) cells labeling MUT with AP22355c at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG (1583138) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm mainly and mitochondrion staining on Hela cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin (PD18466410) at 1/100 dilution (red).The nuclear counter stain is DAPI (blue).



All lanes : Anti-MUT Antibody (Center) at 1:2000 dilution Lane 1: Human liver lysate Lane 2: Mouse liver lysate Lane 3: K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : kDa Blocking/Dilution buffer: 5% NFDM/TBST.

All lanes : Anti-MUT Antibody (Center) at 1:2000 dilution Lane 1: K562 whole cell lysate Lane 2: NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 83 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.