

MAP4K1-T165 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22379a

Product Information

Application	WB, E
Primary Accession	Q92918
Reactivity	Human, Mouse
Predicted	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB59626
Calculated MW	91296

Additional Information

Gene ID	11184
Other Names	Mitogen-activated protein kinase kinase kinase kinase 1, 2.7.11.1, Hematopoietic progenitor kinase, MAPK/ERK kinase kinase kinase 1, MEK kinase kinase 1, MEKKK 1, MAP4K1, HPK1
Target/Specificity	This MAP4K1-T165 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 140-180 from the human region of human MAP4K1-T165.
Dilution	WB~~1:1000-2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MAP4K1-T165 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MAP4K1 (HGNC:6863)
Synonyms	HPK1
Function	Serine/threonine-protein kinase, which plays a role in the response to

environmental stress (PubMed:[24362026](#)). Appears to act upstream of the JUN N-terminal pathway (PubMed:[8824585](#)). Activator of the Hippo signaling pathway which plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. MAP4Ks act in parallel to and are partially redundant with STK3/MST2 and STK4/MST2 in the phosphorylation and activation of LATS1/2, and establish MAP4Ks as components of the expanded Hippo pathway (PubMed:[26437443](#)). May play a role in hematopoietic lineage decisions and growth regulation (PubMed:[24362026](#), PubMed:[8824585](#)). Together with CLNK, it enhances CD3-triggered activation of T-cells and subsequent IL2 production (By similarity).

Tissue Location

Expressed primarily in hematopoietic organs, including bone marrow, spleen and thymus. Also expressed at very low levels in lung, kidney, mammary glands and small intestine


Background

Serine/threonine-protein kinase, which may play a role in the response to environmental stress. Appears to act upstream of the JUN N-terminal pathway. May play a role in hematopoietic lineage decisions and growth regulation. Able to autophosphorylate.

References

Hu M.C.-T.,et al.Genes Dev. 10:2251-2264(1996).
 Grimwood J.,et al.Nature 428:529-535(2004).
 Oppermann F.S.,et al.Mol. Cell. Proteomics 8:1751-1764(2009).
 Mayya V.,et al.Sci. Signal. 2:RA46-RA46(2009).
 Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).

Images



All lanes : Anti-Phospho-MAP4K1-T165 Antibody, ctrl at 1:1000-2000 dilution Lane 1: Jurkat whole cell lysate Lane 2: THP-1 whole cell lysate Lane 3: Ramos whole cell lysate Lane 4: Raji whole cell lysate Lane 5: Daudi whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 91 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.